

Before using, be sure to read this written explanation.
Also, please put it in a safe place so that you can read it when necessary.

Class 2 OTC Medicine For Nasal Allergy Symptoms caused by Pollen, House Dust, etc

AG Allercut® C

Sodium Cromoglicate Nasal Spray

Characteristics of AG Allercut C

1. The anti-allergy component, "sodium cromoglicate", suppresses the release of allergy-inducing substances and relieves severe allergy symptoms.
2. "Chlorpheniramine maleate" blocks the histamine receptor binding, which causes allergy symptoms, and relieves sneezing and runny nose.
3. "Naphazoline hydrochloride" relieves nasal mucosal swelling and congestion, rapidly improving nasal congestion.
4. "Dipotassium glycyrrhizinate" soothes nasal inflammation caused by allergic reactions.
5. "Cetylpyridinium chloride hydrate" suppresses bacterial proliferation with its bactericidal action, keeping the nasal cavity clean.
6. It is a nasal spray with a refreshing and cool sensation.



Precautions



Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects/accidents may occur more readily)

1. **This product should not be used by the following persons.**
Persons who have had allergic symptoms to this product or its ingredients.
2. **Do not drive vehicles or operate machines after using this product.**
(Symptoms such as sleepiness etc. may occur)
3. **Do not use this product continuously for a long term.**



Things to consult

1. **The following persons should consult a doctor, pharmacist, or registered seller before use.**
 - (1) Persons undergoing medical treatment from a doctor
 - (2) Persons undergoing hypo-sensitization (allergy desensitization) therapy or allergy treatments
 - (3) Women who are pregnant or considered to be pregnant
 - (4) Persons who have had allergic symptoms to medicines or such
 - (5) Persons who have been diagnosed with the following conditions:
High blood pressure, heart disease, diabetes, thyroid dysfunction, glaucoma
 - (6) Persons who are unclear whether the symptoms are caused by an allergy or other conditions
2. **If the following symptoms are observed after using this medicine, these may be adverse reactions, so stop using the medicine immediately and bring this instruction to a consultation with your doctor, pharmacist, or registered seller.**

Related Part	Symptoms
Skin	Rash/redness, itching, swelling
Nasal	Swelling, irritation, nosebleeds
Nervous system	Headache

In rare cases, the following serious symptoms may occur. In such cases, seek medical attention immediately.

Symptom Name	Symptoms
Shock (anaphylaxis)	Symptoms such as itchy skin, hives, hoarse voice, sneezing, itchy throat, shortness of breath, palpitations, or impaired consciousness

3. **If symptoms do not improve after taking the product for 3 days, discontinue use and bring this instruction to a consultation with a doctor, pharmacist, or registered seller.**
4. **If using this product for more than 2 weeks, even after symptoms improve, please bring this instruction to a consultation with a doctor, pharmacist, or registered seller.**

Indications

Relief of the following nasal allergy symptoms caused by pollen, house dust (indoor dust), and similar triggers:

Runny nose (excessive nasal discharge), nasal congestion, sneezing, heavy headedness


Dosage and Administration

Age	Amount per dose	Daily dose
Adults (15 years or over)	Spray once into each nostril.	3 to 5 times (With intervals of 3 hours or more)
7 to 15 years		
Under 7 years	Do not use.	

<Precautions related to dosage and administration>

- (1) Follow the Dosage and Administration strictly.
- (2) Excessive use may cause nasal congestion.
- (3) For children aged 7 and older, please follow the guidance of a guardian when using this product.
- (4) Use for nasal use only.

1. Blow your nose thoroughly before use.
(Do not blow too hard.)
2. Remove the blue cap and hold the container as shown in the figure. (At the beginning of use, press several times until the medicinal solution is sprayed.)
3. Gently insert the nozzle tip into the nasal cavity and spray the medicinal solution once. Repeat the same process in the other nostril.
4. After use, wipe the nozzle with a clean tissue paper or similar material, and close the cap.



Ingredients and Quantities

The product is a colorless to light yellow clear solution, containing the following ingredients per 100 mL:

Ingredient	Quantity	Function
Sodium Cromoglicate	1 g	Suppresses the release of allergy-inducing substances and relieves severe allergy symptoms.
Chlorpheniramine Maleate	0.25 g	Blocks histamine receptor binding, which causes allergy symptoms, and relieves sneezing and runny nose.
Naphazoline Hydrochloride	0.025 g	Relieves nasal mucosal swelling and congestion, quickly improving nasal congestion.
Dipotassium Glycyrrhizinate	0.3 g	Suppresses nasal inflammation caused by allergic reactions.
Cetylpyridinium chloride hydrate	0.05 g	Suppresses bacterial proliferation with its bactericidal action, keeping the nasal cavity clean.

Additives: Edetate Sodium, Polysorbate 80, Glycerin, l-Menthol, Fragrance, Ethanol

<Precautions related to ingredients and quantities>

This product contains alcohol and may cause a stinging sensation.

Storage and Handling Precautions

- (1) Ensure the cap is closed then store in a cool place and out of direct sunlight.
- (2) Keep out of reach of children.
- (3) Don't put it in a different container. (It can lead to misuse or lowering of quality.)
- (4) Please do not share this product with others.
- (5) Please do not use this product after the expiration date. Even within the expiration date, please use it promptly after opening.

If you have any questions about this product, please contact the store where you purchased it or at the following address:

Daiichi Sankyo Healthcare Co., Ltd. Customer Relations Office

3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan zip/postal code 103-8234

TEL: 0120-337-336 Reception Hours: 9:00 am - 5:00 pm (closed Saturday, Sunday, and holidays)

<https://www.daiichisankyo-hc.co.jp/>

01® Registered Trademark



Seller
Daiichi Sankyo Healthcare Co., Ltd.
3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan

Manufacturer and Seller
Teika Pharmaceutical Co., Ltd.
1-3-27, Arakawa, Toyama City