

Before using, be sure to read this written explanation.
Also, please put it in a safe place so that you can read it when necessary.

Designated Class 2 OTC
Medicine

For Nasal Congestion, Runny Nose, and Sneezing caused by Pollen

AG Allercut® EXc <Exclusive for Seasonal Allergy>

Beclomethasone Dipropionate (steroid) 0.1% Combination Nasal Spray

Characteristics of AG Allercut EXc <Exclusive for Seasonal Allergy>

1. Beclomethasone dipropionate, which contain both anti-inflammatory and anti-allergic effects, acts directly on the nasal mucosa to soothe nasal congestion, runny nose, and sneezing caused by pollen.
2. This nasal spray remains effective from morning until evening when used in the morning or from evening until morning when used in the evening. (When used twice a day as usual)
3. Provides a refreshing, cool sensation, stays in the affected area and does not drip easily.
4. The spray container is designed to spray effortlessly at any angle until the last drop.
5. It can be used in conjunction with oral medications or eye drops for rhinitis treatment.



Precautions



Things to avoid (If you fail to follow these precautions, your present symptoms may worsen, or side effects may occur more readily.)

1. **This product should not be used by the following persons.**
 - (1) Persons who have been diagnosed with the following conditions:
Systemic mycosis, tuberculosis, high blood pressure, diabetes, recurrent nosebleeds, asthma, glaucoma, infections
 - (2) Persons with purulent nasal passages (accumulation of pus due to hair follicle infection, accompanied by pain and swelling)
 - (3) Persons who have had allergic symptoms to this product or its ingredients
 - (4) Under 18 years old
 - (5) Women who are pregnant or considered to be pregnant
 - (6) Persons who have used steroid nasal sprays for more than 3 months within the past year
2. **Do not use this medication for more than 3 months within a year, including the usage of other steroid nasal sprays. (If usage beyond 3 months is necessary, please consult with an otolaryngologist as there may be other underlying diseases.)**
3. **After using this medication, do not use other steroid nasal sprays. However, if prescribed by a doctor, follow their instructions.**



Things to consult

1. **The following persons should consult a doctor, pharmacist, or registered seller before use.**
 - (1) Persons undergoing medical treatment from a doctor
 - (2) Persons undergoing hypo-sensitization (allergy desensitization) therapy or allergy treatments
 - (3) Persons with pain in the head, forehead, or cheeks, and yellow or green nasal discharge (infectious sinusitis)
 - (4) Persons who are breastfeeding
 - (5) Persons who have had allergic symptoms to medicines or such
 - (6) Persons who are unclear whether the symptoms are caused by a seasonal allergy or other conditions
 - (7) Elderly people
 - (8) Persons with hypertrophic rhinitis or nasal growths^{*1} (nasal polyps) ^{*2}
 - *1: Symptoms include a heavy feeling around the nose and the presence of a small amount of mucus or yellow or green nasal discharge.
 - *2: Symptoms include nasal congestion, nasal voice, and a feeling of something in the back of the nose.
2. **If the following symptoms are observed after using this medicine, these may be adverse reactions, so stop using the medicine immediately and bring this instruction to a consultation with your doctor, pharmacist, or registered seller.**

Related Part	Symptoms
Nasal	Nosebleeds, crusts inside the nose, irritation, itching, dryness, discomfort, paroxysmal sneeze, altered sense of smell, purulent symptoms (accumulation of pus due to hair follicle infection, accompanied by pain and swelling)
Throat	Irritation, foreign body sensation, purulent symptoms (accumulation of whitish pus in the back of the throat due to infection, accompanied by pain and swelling)
Skin	Rash/redness, itching, swelling
Nervous system	Headache, dizziness
Digestive system	Nausea, vomiting, diarrhea, loss of appetite
Others	Onset of asthma, eye pain, blurred vision, palpitations, elevated blood pressure

In rare cases, the following serious symptoms may occur. In such cases, seek medical attention immediately.

Symptom Name	Symptoms
Shock (anaphylaxis)	Symptoms such as itchy skin, hives, hoarse voice, sneezing, itchy throat, shortness of breath, palpitations, or impaired consciousness.

- If pain occurs in the head, forehead, cheeks, or if nasal discharge appears yellow or green, or if unusual symptoms occur after use, so stop using the medicine immediately and bring this instruction on the package to a consultation with your doctor, pharmacist, or registered seller. (Other diseases may underlie.)
- If improvement in symptoms is not observed after approximately 1 week of use (up to 4 times a day, 8 sprays per day), discontinue use and consult a physician, pharmacist, or registered seller with this instruction.

Indications

Relief of symptoms caused by seasonal allergies to pollen: Nasal congestion, runny nose (excessive nasal discharge), sneezing

Dosage and Administration

Age	Amount per dose	Daily dose
Adults (18 years or over)	Spray once into each nostril.	Normally, use twice a day (morning and evening)
Under 18 years	Do not use.	



You can use up to 4 times a day (8 sprays), but make sure to wait at least 3 hours between uses.

- If your symptoms improve, reduce the frequency of use.
- If your symptoms worsen again, you can increase the frequency of use.
- Do not use for more than 3 months within 1 year.

<Precautions related to dosage and administration>

- This product contains beclomethasone dipropionate (steroid). If using it excessively or incorrectly, adverse reactions/incidents are more likely to occur. Please strictly follow the recommended dosage and usage.
- Use for nasal use only.
- If you taste it during use, rinse your mouth.

<How to use>

 <p>1 Nasal Nostrils</p>	 <p>2</p>	 <p>3 Remove the blue cap. Nozzle</p>	 <p>4</p>	 <p>5</p>
Do not use if you have suppuration (accumulation of pus due to hair follicle infection) in your nostrils.	Gently blow your nose before use. (Do not blow too hard)	Remove the blue cap and hold the container as shown in the figure. (At the beginning of use, press several times until the medicinal solution is sprayed)	Insert the tip of the nozzle gently into the nasal cavity and spray the solution once each time. Repeat the same operation in the other nostril.	After use, wipe the area near the nozzle with a clean tissue or the like, and close the cap.

<Precautions for Use>

- Before use, blow your nose and ensure the passage of the nasal cavity is clear.
 - Be careful not to let the container's tip come into contact with nasal discharge or other substances, as it may contaminate the solution.
 - Ensure the nozzle is not directed towards the nasal septum*. Insert it straight into the nasal cavity for spraying. Especially for right-handed persons, the nozzle tends to point towards the right nasal septum, while for left-handed, it tends to point towards the left. Please be cautious.
- *Nasal Septum: The partition that divides the nasal cavities inside the nostrils
- After use, wipe the area near the nozzle with a tissue or similar item to keep it clean.
 - Do not attempt to puncture the tip of the nozzle with a needle or any sharp object, as it can be extremely dangerous if it breaks.

Ingredients and Quantities

The product is a white, suspending viscous liquid containing the following ingredients per 100g.

Ingredient	Quantity	Function
Beclomethasone Dipropionate	0.1 g	Acts on the nasal mucosa to alleviate nasal congestion, runny nose, and similar symptoms.

Additives: Carboxyvinyl Polymer, L-Arginine, Ethanol, Tonicity Agent, Benzalkonium Chloride, Edetate Sodium, Mentha Oil, I-Menthol, Polysorbate 80, pH adjuster

<Precautions related to ingredients and quantities> This product contains alcohol and may cause a stinging sensation.

Storage and Handling Precautions

- (1) Ensure the cap is closed then store in a cool place and out of direct sunlight.
- (2) Keep out of reach of children.
- (3) Don't put it in a different container. (It can lead to misuse or lowering of quality)
- (4) Please do not share this product with others.
- (5) Please do not use this product after the expiration date. Even within the expiration date, please use it promptly after opening.

If you have any questions about this product,
please contact the store where you purchased it or at the following address:
Daiichi Sankyo Healthcare Co., Ltd. Customer Relations Office
3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan zip/postal code 103-8234
TEL: 0120-337-336
Reception Hours: 9:00 am - 5:00 pm (closed Saturday, Sunday, and holidays)

<https://www.daiichisankyo-hc.co.jp/>

03® Registered Trademark



Seller
Daiichi Sankyo Healthcare Co., Ltd.
3-14-10, Nihonbashi, Chuo-ku,
Daiichi-Sankyo Tokyo, Japan

Manufacturer and Seller
Toko Pharmaceutical Industries, Co., Ltd.
Tsuji 20, Tateyamamachi, Nakaniikawa-gun,
Toyama Prefecture