

Before using, be sure to read this written explanation.
Also please put it in a safe place, so that you can read

Antipyretic analgesics

Class 2 OTC medicine

**Quick relief and works well, easy on your stomach
For headache, fever, menstrual pain**

CALONAL® A

Characteristics of CALONAL

- Antipyretic analgesic ingredient [acetaminophen] acts rapidly on the central nervous system and exert an excellent effect on pains and fevers.
- Antipyretic analgesics that are easy on stomach.
- Does not include ingredients that cause drowsiness (Sedative hypnotic ingredients).



Precautions



Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects may occur more readily)

- 1. This product should not be taken by the following persons.**
 - (1) Persons who have had allergic symptoms with this product or its ingredients.
 - (2) Persons who have developed asthma after taking this product, other antipyretic analgesics, or cold medicines.
- 2. Do not take any of the following medicines while taking this product:**
Other antipyretic analgesics, cold medicines, sedatives
- 3. Do not drink alcohol before or after taking this product.**
- 4. Do not use this product continuously for a long term.**



Things to consult

- 1. The following persons should consult a physician, dentist, pharmacist, or registered salesperson before use.**
 - (1) Persons who are receiving treatment from a physician or dentist.
 - (2) Women who are pregnant or supposed to be pregnant
 - (3) Elderly people
 - (4) Persons who have had an allergic reaction to a drug, etc.
 - (5) Persons who have been diagnosed as the following conditions:
heart disease, kidney disease, liver disease, gastric/duodenal ulcer

- 2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this instruction to your physician, pharmacist, or registered salesperson.**

Related Part	Symptoms
Skin	Rash/redness, itching
Digestive system	Nausea/vomiting, loss of appetite
Neuropsychiatric system	Dizziness
Other	Excessive decrease in body temperature

In rare cases, the following serious symptoms may occur. In such cases, seek medical attention immediately.

Symptom Names	Symptoms
Shock (anaphylaxis)	Immediately after taking the medicine, itchy skin, hives, hoarse voice, sneezing, itchy throat, breathlessness, palpitations, and confusion in consciousness may occur.
Mucocutaneous-ocular Syndrome (Stevens-Johnson Syndrome), toxic epidermal necrolysis, acute generalized exanthematous pustulosis	High fever, redness of the eyes, eye discharge, sore lips, sore throat, widespread rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.
* <u>Drug-Induced Hypersensitivity Syndrome (DIHS)</u>	<u>Widespread skin redness, systemic rash, fever, general malaise, swelling of lymph nodes (neck, armpits, groin, etc.).</u>
Impairment of liver function	Fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, general lethargy, and loss of appetite may occur.
Renal impairment	Fever, rash, decreased in urine volume, general swelling, general lethargy, joint pain (pain in the joints), diarrhea, etc.
Interstitial pneumonia	Shortness of breath or breathlessness when climbing stairs or exerting oneself a little, dry cough, fever, etc. These symptoms may appear suddenly or persist.
Asthma	When you breathe, you may hear a wheezing sound or have difficulty breathing.


- 3. If you have taken the medicine for 5-6 times with no sign of symptoms abating, stop taking the medicine and bring this instruction to a consultation with your physician, dentist, pharmacist, or registered salesperson.**

Indications

- Analgesic for headache, menstrual pain, toothache, pain after tooth extraction, sore throat, back pain, joint pain, neuralgia, muscle pain, stiff shoulder pain, earache, bruise pain, fracture pain, sprain pain, trauma pain
- Relief from chills and fever

Dosage and Administration

Please take the following amount with water or lukewarm water.

Age	Amount per dose	Daily dose
Adult (age of 15 and above)	1 tablet 	Take up to 3 times. Please avoid taking it on an empty stomach if possible. The interval between doses should be at least 4 hours.
Under 15 years	Do not use.	

<Precautions related to Dosage and Administration>

- (1) Strictly follow the dosage and administration guidelines.
- (2) How to take out the tablet

As shown in the illustration on the right, push the tablet out firmly with the fingertip from the convex side of the PTP sheet through the aluminum foil on the other side to put out the tablet. (If the PTP sheet is swallowed, the sharp corners of the sheet may puncture the esophageal mucosa or other accidents may occur.)



Ingredients and Quantities

This product is a white, uncoated tablet containing the following ingredients per 1 tablet.

Ingredients	Amount	Function
Acetaminophen	300 mg	Acts on the central nervous system, reduces fever and relieves pains.

Additives: Cellulose, Partly Pregelatinized Starch, Partially Hydrolyzed Polyvinyl Alcohol, Magnesium Stearate, l-Menthol, Fragrance, Lactose

Storage and Handling Precautions

- (1) Store in a cool place with little humidity and out of direct sunlight.
- (2) Store out of reach of children.
- (3) Do not put in a different container. (It can lead to misuse or lowering of quality.)
- (4) Do not use the product after the indicated expiration date.

If you have any questions about this product, please contact the store where you purchased it or at the following address:
Daiichi Sankyo Healthcare Co., Ltd. Customer Relations Office
3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan zip/postal code 103-8234
TEL: 0120-337-336
Reception Hours: 9:00 am -5:00 pm (closed Saturday, Sunday, and holidays)

<https://www.daiichisankyo-hc.co.jp/>

02® Registered Trademark



Manufacturer and seller
Daiichi Sankyo Healthcare Co., Ltd.
3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan