

Stimulates the stomach functions and improves the intestinal condition to its normal state

MADE IN JAPAN

第一三共胃腸薬プラス 細粒

Daiichi Sankyo Ichoyaku Plus Fine Granules

Digestive medicine

Class 2 OTC medicine



Indications

- Stomach upset, overeating, excessive drinking, pressure on the chest, loss of appetite
- Heartburn, stomachache, chlorhydria, stomach heaviness, stomach discomfort, belching
- Dyspepsia, accelerated digestion, indigestion, a feeling of gastric/abdominal distension
- Nausea (heartburn, nausea due to hangover/ drunken sickness), vomiting
- Intestinal regulation (regulating bowel movement), soft stool, constipation



Dosage and Administration

Take the following amount with cold or warm water.

Age	Amount per dose	Daily dose
15 years or over	1 sachet	3 times Take after meals
11 to 14 years	2/3 sachet	
8 to 10 years	1/2 sachet	
5 to 7 years	1/3 sachet	
3 to 4 years	1/4 sachet	
Under 3 years	Do not use.	

<Precautions related to Dosage and Administration>

- (1) Follow the Dosage and Administration strictly.
- (2) Use for infants aged 3 or over should be supervised by guardians.

Ingredients and Quantities

This product is a sachet containing white to slightly yellow-brownish white fine granules and pale brown to brown fine granules, and has a specific smell. Three sachets (1.3 g per sachet) contain the following ingredients.

Ingredient	Quantity
Taka-diastrase N1	150 mg
Lipase AP12	60 mg
Lactobacillus sporegene (bulk powder of Lacbon)	60 mg
Magnesium aluminosilicate	900 mg
Synthetic hydrotalcite	600 mg
Precipitated calcium carbonate	600 mg
Powdered phellodendron bark	105 mg
Powdered cinnamon bark	225 mg
Powdered ginger	75 mg
Powdered clove	30 mg
Powdered fennel	60 mg
<i>l</i> -menthol	9 mg
Aldioxa	60 mg
Powdered glycyrrhiza	150 mg

Other ingredients: Cellulose, lactose, polysorbate 80, hydroxypropyl cellulose, Japanese pepper

Precautions

Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects may occur more readily.)

1. This product should not be used by the following persons:
Persons who are getting dialysis therapy.
2. Do not use this product continuously for a long term.

Things to consult

1. The following persons should consult a doctor or pharmacist before use.
 - (1) Persons who are receiving medical treatment from a doctor.
 - (2) Persons who have had allergic symptoms with medicines or such.
 - (3) Persons who have been diagnosed as the following conditions:
Kidney disease, thyroidal functional disorder