

Before using, be sure to read this written explanation on the package. Also, please put it in a safe place so that you can read it when necessary.

Anti-inflammatory analgesic Class 2 OTC medicine **7 sheets**
LOXONIN[®]S Tape (7 sheets x 1 bag)

Contains [Ibuprofen sodium hydrate] with excellent anti-inflammatory and analgesic effects.

Characteristic

Activates when absorbed through skin and penetrates directly to the core of severe pain.

Apply once per day. Comfortable-fit tape has excellent stretchability making it less likely to fall off.



Precautions



Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects/accidents may occur more readily.)

1. This product should not be used by the following persons:

- (1) Persons who have experienced allergic symptoms to this product or the ingredients contained in it in the past
- (2) Persons who have experienced asthma after using this product or other antipyretic analgesics, cold medicines, or topical anti-inflammatory analgesics
- (3) Children under the age of 15

2. Please avoid applying to the following parts.

- (1) Around eyes, mucous membranes
 - (2) Skin with rashes, skin irritations, or wounds
 - (3) Athlete's foot, ringworm infections, or other infected areas
3. Please do not use other topical anti-inflammatory analgesics while using this product.

4. Do not use this product continuously for more than 2 weeks.

(This product is intended to temporarily relieve pain. If pain or other symptoms persist, stop using it, and consult a doctor.)



Things to consult

1. The following persons should consult a doctor, pharmacist, or registered seller before use.

- (1) Persons undergoing medical treatment from a doctor
- (2) Persons who have had allergic symptoms to medicines or such
- (3) Women who are pregnant or are considered to be pregnant.
- (4) Elderly people
- (5) Persons with the following diagnoses: Bronchial asthma

2. If the following symptoms are observed after using this product, these may be adverse reactions, so stop using the medicine immediately and bring this instruction on the package to a consultation with your doctor, pharmacist, or registered seller.

Related Part	Symptoms
Skin	Rash, redness, itching, Swelling, tingling, contact dermatitis, blisters, blue bruise, hyperpigmentation
Digestive organs	Stomach discomfort, pain in the solar plexus
Other	Oedema

In rare cases, the following severe symptoms may occur. If they do occur, immediately consult with your doctor.

Symptom Names	Symptoms
Shock (anaphylaxis)	Symptoms such as itchy skin, hives, hoarse voice, sneezing, itchy throat, shortness of breath, palpitations, or impaired consciousness

3. The following symptoms may appear, but if any of these symptoms persist or increase, discontinue use and bring this instruction to a consultation with a doctor, pharmacist, or registered seller.

Diarrhea, loose stool

4. If symptoms do not improve after taking the product for 5 or 6 days, discontinue use and bring this instruction to a consultation with a doctor, pharmacist, or registered seller. (There may be other potential diseases)

Indications

Lower-back pain, shoulder pain associated with stiff shoulders, joint pain, muscle pain, tendonitis (hand/wrist pains), elbow pain (tennis elbow, etc.), bruises, sprains

Dosage and Administration

Peel off the surface liner (film) and apply to the affected area once a day.

<Precautions related to dosage and administration> (1) Follow the dosage and administration strictly. (2) This product is intended to relieve symptoms such as pain or swelling. It is not intended to cure the causes of pain or swellings. Therefore, apply this product only when symptoms appear. (3) Do not use more than 4 sheets per day. (4) If you are sweating or the affected area gets wet, dry it off thoroughly before applying. (5) For persons with sensitive skin, cut the sheet into a 1-2 cm square and apply it to a delicate area such as the inner arm for at least half a day. Confirm that no symptoms such as rashes, redness, itching, or irritations appear before use.

Ingredients and Quantities (per 100g plaster)

Ingredient	Quantity	Effect
Loxoprofen sodium hydrate	5.67g (5g as anhydride)	Suppresses [prostaglandin], the source of inflammation and pain.

Additives: Styrene-isoprene-styrene-block copolymer (SIS), polyisobutylene, hydrogenated rosin glycerol ester, BHT, l-menthol, liquid paraffin, and 2 other ingredients

[1g plaster per sheet (7cm×10cm)]

Storage and Handling Precautions

(1) Store in a cool place with little humidity and out of direct sunlight. (2) Keep out of reach of children. (3) Don't put it in a different container. (It can lead to misuse or a lowering of quality.) (4) To maintain product quality, store unused sheets in the bag and seal it properly. (5) Please do not use medicine after the expiration date. It is recommended to use the product shortly after opening an aluminum bag.

If you have any questions about this product, please contact the store where you purchased it or at the following address:
Daiichi Sankyo Healthcare Co., Ltd. Customer Relations Office
3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan zip/postal code 103-8234
TEL: 0120-337-336 Reception Hours: 9:00 am -5:00 pm
(Closed Saturday, Sunday, and holidays)

<https://www.daiichisankyo-hc.co.jp/>
07@registered trademark

Relief System for Sufferers from
Adverse Drug Reactions
TEL0120 - 149 - 931

Seller

Daiichi Sankyo Healthcare Co., Ltd.

3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan

Manufacturer and seller

Lead Chemical Co., Ltd.

77-3, Himata, Toyama-City, Toyama, Japan

Made in Japan