マキロシ かゆみどめ液P

Makiron Anti-itching Lotion P

Antipruritic lotion

Class 3 OTC medicine



Indications

Insect bites, itchiness

Dosage and Administration

Apply an appropriate amount several times a day onto the affected area.

- <Precautions related to Dosage and Administration>
- (1) Follow the Dosage and Administration strictly.
- (2) Use for children should be supervised by guardians.
- (3) Be careful about the intrusion of medicinal liquid into the eyes. If it happens, rinse your eyes immediately with water or lukewarm water. If symptoms are severe, see an ophthalmologist.
- (4) This product is for external use only.

<<How to apply>>

Hold the bottle upside down and press the application rubber onto the affected area several times to immerse the rubber with the medicinal liquid adequately. Note that the rubber may be broken if it is not adequately immersed with the medicinal liquid. Be sure to close the cap tightly to prevent the rubber from drying after use. The rubber may be deteriorated or broken by the adhesion of oily materials such as cream or ointment.



Ingredients and Quantities

In 100 mL,

Ingredient	Quantity
Diphenhydramine hydrochloride	2.0 g
Glycyrrhetinic acid	0.2 g
Isopropyl methylphenol	0.1 g
<i>I</i> -menthol	5.0 g
dl-camphor	0.1 g

Other ingredients: 1,3-butylene glycol, citric acid, sodium citrate, hydroxypropyl cellulose, ethanol As this product contains alcohol, it may stimulate the applied area.

Precautions

Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects may occur more readily.)

Do not use at the following sites:

- (1) Around the eyes, mucous membranes or such
- (2) Affected site with a wound

Things to consult

- 1. The following persons should consult a doctor or pharmacist before use.
 - (1) Persons who are receiving medical treatment from a doctor.
 - (2) Persons who have had allergic symptoms with medicines or such.
 - (3) Persons who have severe wetting or erosion of skin.