

Inconspicuous transparent patch which avoid scratching

MADE IN JAPAN

# マキロン®パッチエース

## Makiron Patches Ace

Antipruritic and antiphlogistic patch

Designated class 2 OTC medicine



### Indications

Itching due to insect bites

### Dosage and Administration

Apply to the affected area 1 to 3 times a day.

<Precautions related to Dosage and Administration>

- (1) Follow the Dosage and Administration strictly.
- (2) Use for children should be supervised by guardians.
- (3) This product is for external use only.
- (4) When the affected area is sweating or wet, wipe away sufficiently before use.
- (5) If itching resolves after using this product for a few hours, please remove it.
- (6) Do not use at the same site for a long time.

### Ingredients and Quantities

In 100 g (1.25 m<sup>2</sup>) of plaster of this product:

Ingredients	Quantity
Dexamethasone acetate	0.025 g
Diphenhydramine	1.00 g
Isopropyl methylphenol	1.00 g
Tocopherol acetate (acetate ester of vitamin E)	0.50 g

Additives: Diisopropyl adipate, methacrylate/ butylacrylate copolymer, BHT, *l*- menthol

This product contains dexamethasone acetate, an adrenocortical hormone (steroid).

## Precautions

### Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects may occur more readily.)

1. Do not use at the following sites:
  - (1) Varicella (chickenpox), athlete's foot, ringworm or suppurative area
  - (2) Area around the eyes, mucous membrane of the eyes and such
  - (3) Wound, moist skin (scratch), erosion, eczema, eruption
2. This product should not be used on large areas of the face.
3. Do not use this product continuously for a long term.

### Things to consult

1. The following persons should consult a doctor or pharmacist before use.
  - (1) Persons who are receiving medical treatment from a doctor.
  - (2) Women who are pregnant or considered to be pregnant.
  - (3) Persons who have had allergic symptoms with medicines or such.
  - (4) Persons whose affected areas are large.