

Before using, be sure to read these written explanation.

Also please put it in a safe place, so that you can read them when necessary.

When you have a bad cold, take in the morning and it keeps on working until night.

# プレコール® CR 持続性錠

## Precol® CR Long-lasting Tablets


Common cold medicine <Pyrene medicine>



Designated Class 2 OTC Medicine

### Characteristics of the Precol CR Long-lasting Tablets

- Seven active ingredients, including two antipyretic analgesic ingredients (isopropylantipyrine, acetaminophen) produce an effective, balanced attack against sore throats, fever, and other symptoms of a bad cold.
- Our original formulation technology (Control Release Technology) features two layers (the fast-dissolving and long-lasting layers) that release the medicine into your body differently.
- Take this easy-to-swallow tablet common cold medicine twice a day, in the morning and at night.



### Indications

Relief of various symptoms of a common cold (sore throat, fever, runny nose, stuffy nose, cough, sputum, chills, headache, joint pain, muscle pain, and sneezing)

### Administration and Dosage

Take the following amount with cold or warm water.

Age	Amount per dose	Daily dose
Adults (15 years or over)	3 tablets	twice (Morning and night, preferably within 30 minutes after your meal)
Under 15 years	Do not use.	

<Precautions related to Dosage and Administration>

- (1) Follow the Dosage and Administration strictly.
- (2) How to take out the tablets:

As shown in the diagram, use your finger to push the PTP sheet that contains the tablet. Take off the aluminum foil on the back and ingest the tablet. (If you mistakenly take the tablet with the wrapping on it, it could puncture the mucous membrane of the esophagus and lead to other unknown consequences.)



## Ingredients and Quantities

This product is white film-coated tablet. Six tablets contain the following ingredients.

Ingredient	Quantity	Function
Isopropylantipyrine (pyrene medicine)	300mg	Reduces fever, relieves sore throat, headache, and joint pain.
Acetaminophen	450mg	
Chlorpheniramine maleate	6mg	Relieves runny nose, stuffy nose, sneezing.
Dihydrocodeine phosphate	18mg	Relieve coughs.
dl-methylephedrine hydrochloride	60mg	Relieve coughs and reduces phlegm.
Powdered extract of glycyrrhiza	90mg (750mg as crude drug)	Eases inflammation of the mucous membranes in the nose and throat.
Anhydrous caffeine	75mg	Eases headaches.

Additives: lactose, hydroxypropyl cellulose, cellulose, Magnesium stearate, hypromellose, methacrylic acid copolymer L, titanium oxide, macrogol, talc, stearic acid, carnauba wax.

## Precautions

### Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects/accidents may occur more readily.)

- This product should not be used by the following persons:
  - Persons who have had allergic symptoms with this product or its ingredients.
  - Persons who have developed asthma after taking this product, other cold medicines or antipyretic analgesics.
  - Children younger than 15.
- Do not use any of the following medicines while taking this product:  
Other cold medicines, antipyretics, analgesics, sedatives, antitussives, expectorants, oral medicines containing antihistamine (oral medicines for rhinitis, motion sickness medicines, anti-allergic medicines or such)
- Do not drive vehicles or operate machines after taking this product (Sleepiness or such may occur).
- Women who are breastfeeding should not take this product or should avoid breastfeeding if take it.
- Do not drink alcohol before or after taking this product.
- Do not use this product continuously for a long term.

### Things to consult

- The following persons should consult a doctor or pharmacist before use.
  - Persons who are treated by a doctor or dentist
  - Women who are pregnant or considered to be pregnant
  - Elderly people
  - Persons who have had allergic symptoms with medicines or such.
  - Persons who have the following symptoms: High fever, dysuria
  - Persons who have been diagnosed as the following conditions:  
Thyroidal functional disorder, diabetes, heart disease, hypertension, liver disease, kidney disease, gastric/duodenal ulcer, glaucoma, respiratory functional disorder, obstructive sleep apnea syndrome, obesity

## Precautions for storage and handling

- (1) Please store in a cool and dry place away from sunlight.
- (2) Please keep out of the reach of children.
- (3) Please do not transfer this product to a different container. (It may cause misuse and affect the quality.)
- (4) Please do not use after the expiration date printed on the label.

Please direct questions about this product to the store where you purchased it or to the below address:

Customer Relations Center  
Dai-Ichi Sankyo Healthcare Co., Ltd.  
3-14-10, Nihonbashi, Chuo-ku, Tokyo, 103-8234, Japan  
TEL: 0120-337-336  
Operation Hours: 9am-5pm (not including Sat., Sun. and holidays)

<https://www.daiichisankyo-hc.co.jp/>

01@Registered trademark

Manufactured and Sold by  
Dai-Ichi Sankyo Healthcare Co., Ltd.  
3-14-10, Nihonbashi, Chuo-ku, Tokyo

英語 (English) 2020.7 作成

This document is reference information about the products. Please confirm a document attached to a product definitely.