

For eye allergy symptoms caused by pollen, house dust or such.

MADE IN JAPAN

エージーズ® **アレルギーカット[®]C**

AG Eyes Allercut C

Anti-allergy eye drops containing sodium cromoglicate

Class 2 OTC medicine



Indications

Relief of the following eye allergy symptoms caused by pollen, house dust, or such: Itchy eye, red eye, blurred vision (e.g., when having a lot of eye mucus), tearful eyes and sensation of a foreign object in the eye (gritty sensation)

Dosage and Administration

Put 1 to 2 drops per dose, 4 to 6 times a day.

<Precautions related to Dosage and Administration>

- (1) Follow the Dosage and Administration strictly.
- (2) Use for children should be supervised by guardians.
- (3) Keep the tip of dropper away from the eyelid or the eyelashes.
Do not use the product if the medicinal liquid has been cloudy.
- (4) Do not use this product while you are wearing contact lenses.
- (5) This product is for ocular use only.

1. Wash hands elaborately before use.
2. Pull down the lower lid of the eye lightly. Then drop off 1 to 2 drops of medicinal liquid to the eye. At that time, keep the tip of dropper away from the eyelid or eyelashes.
3. Close the eye, and let the medicinal liquid be absorbed after its application.
4. After use, keep the tip and cap of dropper clean, and close the cap tightly.



Ingredients and Quantities

This product is colorless to pale yellow clear eye drops and contains the following ingredients per 100 mL.

Ingredient	Quantity
Sodium cromoglicate	1 g
Chlorpheniramine maleate	0.015 g
Dipotassium glycyrrhizinate	0.125 g
Sodium chondroitin sulfate	0.2 g

Other ingredients: Sodium edetate, boric acid, borax, benzalkonium chloride, propylene glycol, polysorbate 80, *dl*-camphor, *d*-borneol, pH adjuster, sodium hyaluronate

Precautions

Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects/accidents may occur more readily.)

1. This product should not be used by the following persons:

Persons who have had allergic symptoms with this product or its ingredients.

2. When a nasal spray is used in combination, do not drive vehicles or operate machines after using this product. (Sleepiness or such may occur.)

Things to consult

1. The following persons should consult a doctor or pharmacist before use:

(1) Persons who are receiving medical treatment from a doctor.

(2) Persons who are receiving medical treatment for allergy such as desensitization therapy.

(3) Women who are pregnant or considered to be pregnant.

(4) Persons who have had allergic symptoms with medicines or such.

(5) Persons who have the following symptom: Intense pain in an eye

(6) Persons who have been diagnosed as the following conditions: Glaucoma

(7) Persons who can not assure the reason of their symptoms as an allergy.

Particularly, as the following cases can not be attributed to allergy, consult a doctor before use:

- When symptoms appear only in one eye.
- When having symptoms not in the nose but only in the eyes.
- When the eyesight is also affected.