

Quickly dissolves and acts on upset and discomfort of stomach

MADE IN JAPAN

第一三共胃腸薬〔細粒〕a

Daiichi Sankyo Ichoyaku [Fine Granules] a

Digestive medicine

Class 2 OTC medicine



Indications

- Stomach upset, overeating, excessive drinking, pressure on the chest, loss of appetite
- Heartburn, stomachache, chlorhydria, stomach heaviness, stomach discomfort, belching
- Dyspepsia, accelerated digestion, indigestion, a feeling of gastric /abdominal distension
- Nausea (heartburn, nausea due to hangover/ drunken sickness), vomiting



Dosage and Administration

Take the following amount with cold or warm water.

Age	Amount per dose	Daily dose
15 years or over	1 sachet	3 times Take after meals
11 to 14 years	2/3 sachet	
8 to 10 years	1/2 sachet	
5 to 7 years	1/3 sachet	
3 to 4 years	1/4 sachet	
Under 3 years	Do not use.	

<Precautions related to Dosage and Administration>

- (1) Follow the Dosage and Administration strictly.
- (2) Use for infants aged 3 or over should be supervised by guardians.

Ingredients and Quantities

Three sachets (1.3 g per sachet) of this product contain the following ingredients.

Ingredient	Quantity
Taka-diastase N1	150 mg
Lipase AP12	60 mg
Mallotus japonicus extract	63 mg (504 mg as Mallotus japonicus)
Powdered glycyrrhiza	150 mg
Magnesium aluminosilicate	1200 mg
Synthetic hydrotalcite	450 mg
Magnesium hydroxide	600 mg
Scopolia extract	30 mg
Powdered phellodendron bark	105 mg
Powdered cinnamon bark	225 mg
Powdered fennel	60 mg
Powdered clove	30 mg
Powdered ginger	75 mg
<i>l</i> -menthol	9 mg

Other ingredients: Cellulose, lactose, polysorbate 80, hydroxypropyl cellulose, Japanese pepper

Precautions

Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects may occur more readily.)

1. This product should not be used by the following persons:
Persons who are getting dialysis therapy.
2. Do not use the following medicines while taking this product:
Gastrointestinal analgesic/ antispasmodic medicines
3. Women who are breastfeeding should not take this product or should avoid breastfeeding if take it.
(The ingredients may enter into breast milk, and the baby's heartbeat may be increased.)
4. Do not use this product continuously for a long term.

Things to consult

1. The following persons should consult a doctor or pharmacist before use.
 - (1) Persons who are receiving medical treatment from a doctor.
 - (2) Women who are pregnant or considered to be pregnant.
 - (3) Elderly people.
 - (4) Persons who have had allergic symptoms with medicines or such.
 - (5) Persons who have the following symptom:
Dysuria
 - (6) Persons who have been diagnosed as the following conditions:
Kidney disease, heart disease, glaucoma