

カロヤンプログレ[®]EX O

Karoyan Progre EX O

Hair growth stimulant



Class 3 OTC medicine

Indications

- Premature alopecia, alopecia areata, diffuse alopecia, alopecia pityroides
- Hair growth stimulation, hair restoration, prevention of hair loss (epilation), thinning hair
- Dandruff, itching
- Loss of hair after illness/ childbirth



Dosage and Administration

Age	Amount per dose	Daily dose
Adults (15 years or over)	Rub 2mL onto the scalp, and mildly massage the scalp. Decrease the quantity as necessary if 2mL is too much according to the condition of the affected site.	Twice (morning and evening)
Under 15 years	Do not use.	

<Precautions related to Dosage and Administration>

- (1) Follow the Dosage and Administration strictly. Do not exceed the directed amount and frequency as it may increase the possibility of side effects.
- (2) When using just after hair washing or bathing, cool down the hair and body before use since the side effects such as sweating are tend to occur.
- (3) As elderly persons generally have decreased physiological function, use this product carefully including reduction of the amount per dose or such.
- (4) Be careful to prevent the medicinal liquid from entering the eyes. If it happens, rinse the eyes immediately with water or lukewarm water. If symptoms are severe, see an ophthalmologist.
- (5) To avoid the irritation to mucous membranes of eyes or such by the medicinal liquid left in hands, wash hands elaborately after applying this product.
- (6) This product should be used only on the scalp.

Ingredients and Quantities

This product is a yellow-brown solution, has a slightly specific smell, and contains the following ingredients per 100 mL.

Ingredient	Quantity
Carpronium chloride hydrate	2.18 g (2 g as carpronium chloride)
Panax rhizome tincture	3 mL (1 g as raw crude drug)
Polygonum multiflorum tincture	3 mL (1 g as raw crude drug)
Pyridoxine hydrochloride	0.03 g
Hinokitiol	0.05 g
Pantothenyl ethylether	1 g
<i>l</i> -menthol	0.3 g

Other ingredients: Hydroxypropylchitosan solution, ethanol, pH adjuster, Yellow No. 5

Precautions

Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects may occur more readily.)

Do not use at the following sites:

- (1) Scalp with wound, eczema or inflammation (reddening) or such
- (2) Areas other than the scalp

Things to consult

1. The following persons should consult a doctor or pharmacist before use.
 - (1) Persons who have had allergic symptoms with medicines or cosmetics.
 - (2) Elderly people (See the Section "Precautions related to Dosage and Administration.")