

Before using, be sure to read this written explanation.
Also please put it in a safe place, so that you can read them when necessary.

01

Common cold medicine

Pharmaceuticals requiring guidance

LOXONIN Common cold medicine



Precautions



Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects/accidents may occur more readily)

1. This product should not be taken by the following persons.

- (1) Persons who have had allergic symptoms with this product or its ingredients.
- (2) Persons who have developed asthma after taking this product, other cold medicines, or antipyretic analgesics.
- (3) Children under 15 years old
- (4) Persons undergoing treatment for the following conditions at medical institutions:
Stomach / duodenal ulcers, liver disease, kidney disease, heart disease
- (5) Persons who have been diagnosed by a doctor with the blood abnormalities (blood diseases) such as low red blood cell count (anemia), low platelet count (difficulty in stopping bleeding, bleeds easily), and low white blood cell count etc.
- (6) Expectant women within 12 weeks of delivery date.

2. Do not use any of the following medicines while taking this product:

Other cold medicines, antipyretic analgesics, sedatives, antitussives, expectorants, oral medicines containing antihistamine (oral medicines for rhinitis, motion sickness medicines, anti-allergic medicines or such)

3. Do not drive vehicles or operate machines after taking this product (Drowsiness and other symptoms may occur.)

4. Women who are breastfeeding should not take this product or should avoid breastfeeding if take it.

5. Do not drink alcohol before or after taking this product.

6. Do not take this product for more than 5 days.



Things to consult

1. The following persons should consult a physician or pharmacist.

- (1) Persons who are receiving treatment from a physician or dentist.
- (2) Women who are pregnant or supposed to be pregnant
- (3) Elderly people
- (4) Persons who have had an allergic reaction to a drug, etc.
- (5) Persons who have the following symptoms: High fever, dysuria
- (6) Persons who have been diagnosed as the following conditions:
Thyroidal functional disorder, diabetes, heart disease, hypertension, glaucoma, bronchial asthma, ulcerative colitis, Crohn's disease, lupus erythematosus, mixed connective tissue disease, respiratory dysfunction, obstructive sleep apnea, obesity
- (7) Persons who experienced any of the following diseases:
Stomach / duodenal ulcers, liver disease, kidney disease, blood disease

2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this instruction to your doctor or pharmacist.

- (1) Symptoms such as excessive decrease in body temperature, weakness (lack of strength), and cooling of the extremities (cold hands and feet) are observed after taking this medicine.
- (2) If you experience peptic ulcers or swelling after taking this medicine.
In addition, in rare cases, the following severe symptoms such as gastrointestinal bleeding (vomiting blood, nausea and vomiting, abdominal pain, black tarry stools, bloody stools, etc.), gastrointestinal perforation (perforation of the gastrointestinal tract; nausea, vomiting, severe abdominal pain, etc.), and narrowing or obstruction of the small or large intestine (nausea, vomiting, abdominal pain, abdominal distention, etc.) may occur. In such cases, seek medical attention immediately.

(3) If the following symptoms occurs after taking this medicine

Related Part	Symptoms
Skin	Rash/redness, itching
Digestive system	Abdominal pain, gastric discomfort, loss of appetite, nausea and vomiting, abdominal distention, heartburn, mouth ulcers, indigestion
Cardiovascular system	Increased blood pressure, palpitations
Neuropsychiatric system	Numbness, dizziness, headache, excitement, convulsions
Urinary system	Urination difficulty, hematuria
Other	Chest pain, fatigue, facial hot flashes, fever, anemia

In rare cases, the following serious symptoms may occur. In such cases, seek medical attention immediately.

Symptom Names	Symptoms
Shock (anaphylaxis)	Immediately after taking the medicine, itchy skin, hives, hoarse voice, sneezing, itchy throat, breathlessness, palpitations, and confusion in consciousness may occur.
Blood disorders	Sore throat, fever, general lethargy, pale face and whitish underneath the eyelids, easy bleeding (bleeding gums, nose bleeds, etc.), blue bruises (color does not disappear when pressed), etc.
Mucocutaneous-ocular Syndrome (Stevens-Johnson Syndrome) toxic epidermal necrolysis, erythema multiforme, acute generalized exanthematous pustulosis	High fever, redness of the eyes, eye discharge, sore lips, sore throat, widespread rash/redness, blisters on red areas of the skin, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.
Renal impairment	Fever, rash, decreased in urine volume, general swelling, general lethargy, joint pain (pain in the joints), diarrhea, etc.
Congestive heart failure	General lethargy, palpitations, shortness of breath, chest discomfort, chest pain, dizziness, fainting, etc.
Interstitial pneumonia	Shortness of breath or breathlessness when climbing stairs or exerting oneself a little, dry cough, fever, etc. These symptoms may appear suddenly or persist.
Impairment of liver function	Fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, general lethargy, and loss of appetite may occur.
Rhabdomyolysis	Muscle aches in limbs, shoulders, and hips; numbness, weakness, and stiffness in limbs, general lethargy, and reddish-brown urine.
Aseptic meningitis	Severe headache with neck tightness, fever, nausea and vomiting may occur. (These symptoms have been reported in many patients, especially those undergoing treatment for systemic lupus erythematosus or mixed connective tissue disease.)
Asthma	When you breathe, you may hear a wheezing sound or have difficulty breathing.
Hypoventilation	Symptoms, such as shortness of breath and breathing difficulties, etc. may occur.

3. If any of the following symptoms appear, and persist or increase after taking this medicine, discontinue taking the product and consult a doctor, or pharmacist with this instruction.

Dry mouth, constipation, diarrhea, drowsiness


4. If you have taken the medicine for 5-6 times with no sign of symptoms abating, stop taking the medicine and bring this instruction to a consultation with your doctor or pharmacist. (Especially if a fever lasts more than 3 days or recurring)

Indications

Relief of various symptoms of a cold (runny nose, nasal congestion, sneezing, sore throat, cough, phlegm, chills, fever, headache, joint pain, muscle pain)

Dosage and Administration

Take the following amount with cold or warm water.

Age	Amount per dose	Daily dose
Adult (age of 15 and above)	2 tablets 	3 times Take within 30 minutes after a meal if possible.
Under 15 years	Do not use.	

<Precautions related to Dosage and Administration>

(1) Strictly follow the dosage and administration guidelines.

(2) How to take out the tablet

As shown in the illustration on the right, push the tablet out firmly with the fingertip from the convex side of the PTP sheet through the aluminum foil on the other side to put out the tablet. (If the PTP sheet is swallowed, the sharp corners of the sheet may puncture the esophageal mucosa or other accidents may occur.)



Ingredients and Quantities

This product is white film coated tablet containing the following ingredients per 6 tablets.

Ingredients	Amount	Function
Loxoprofen sodium hydrate	204.3 mg (180 mg as anhydrate)	Reduces fever and relieves sore throat by acting on 【prostaglandins】, the cause of inflammation and pain.
Dihydrocodeine phosphate	24 mg	Acts on the coughing center to relieve coughing.
dl-methylephedrine hydrochloride	60 mg	Expands bronchial tubes and relieves coughing.
Bromhexine hydrochloride	12 mg	Facilitates the transform of phlegm, one of the causes of coughing.
Clemastine fumarate	1.34 mg	Persistently suppresses allergic symptoms of a cold (runny nose, nasal congestion, etc.).

Additives: D-Mannitol, Cellulose, Hydroxypropylcellulose, Croscarmellose Sodium, Magnesium Stearate, Hypromellose, Titanium oxide, Talc, Carnauba Wax

Storage and Handling Precautions

- (1) Store in a cool place with little humidity and out of direct sunlight.
- (2) Store out of reach of children.
- (3) Do not put in a different container. (It can lead to misuse or lowering of quality.)
- (4) Do not use the product after the indicated expiration date.

If you have any questions about this product, please contact the store where you purchased it or at the following address:
Daiichi Sankyo Healthcare Co., Ltd. Customer Relations Office
3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan zip/postal code 103-8234
TEL: 0120-337-336
Reception Hours: 9:00 am -5:00 pm (closed Saturday, Sunday, and holidays)

<https://www.daiichisankyo-hc.co.jp/>

01



Manufacturer and seller
Daiichi Sankyo Healthcare Co., Ltd.
3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan