Before using, be sure to read this written explanation on the package. Also, please put it in a safe place so that you can read it when necessary.

Anti-inflammatory analgesic Class 2 OTC medicine 7 sheets

LOXONIN₈S Tape with Warm Sensation

(7 sheets x 1 bag)

Char acter istic


Contains [loxoprofen sodium hydrate] with excellent anti-inflammatory & analgesic effects.

Warming comfort ingredients are added to the formula. The pleasant warming comfort effect promotes blood circulation in the affected area and relieves pain.

Apply once per day Comfortable-fit tape has excellent stretchability making it less likely to fall off.



Precautions



(If you do not follow these instructions, the current symptoms may worsen, or adverse reactions/incidents are more likely to occur.)

- 1. This product should not be used by the following persons.
- (1) Persons who have experienced allergic symptoms to this product or the ingredients contained in it in the past.
- (2) Persons who have experienced asthma after using this product or other antipyretic analgesics, cold medicines, or topical anti-inflammatory analgesics.
- (3) Children under the age of 15
- 2. Please avoid applying to the following parts.
 - (1) Around eyes, mucous membranes
 - (2) Skin with rashes, skin irritations, or wounds
- (3) Athlete's foot, ringworm infections, or other infected areas
- 3. Please do not use other topical anti-inflammatory analgesics while using this product.
- 4. Do not use this product continuously for more than 2 weeks. (This product is intended to temporarily relieve pain. If pain or other symptoms persist, stop using it and consult a doctor)

Things to consult

- 1. If you fall into any of the following categories, please consult a doctor before use.
- (1) Persons undergoing medical treatment from a doctor.
- (2) Persons who have had allergic symptoms to medicines or such.
- (3) Women who are pregnant or are considered to be pregnant.
- (4) Elderly people
- (5) Persons with the following diagnoses: Bronchial asthma

2. If the following symptoms appear after using this product, you could be experiencing side effects. In such cases, discontinue use immediately and bring this instruction to a consultation with a doctor, pharmacist, or registered seller.

Related Part	Symptoms		
Skin	Rash, redness, itching, swelling, pains, tingling, contact dermatitis, blisters, blue bruises, hyperpigmentation		
Digestive	Stomach discomfort, pain in the solar plexus		
organs			
Other	Oedema		

In rare cases, the following severe symptoms may occur. If they do occur, immediately consult with your doctor.

Symptom Nam	es Symptoms
Shock (anaphylaxis)	Symptoms such as itchy skin, hives, hoarse voice, sneezing, itchy throat, shortness of breath, palpitations, or impaired consciousness

3. The following symptoms may appear, but if any of these symptoms persist or increase, discontinue use and bring this instruction to a consultation with a doctor, pharmacist, or registered seller.

Diarrhea, loose stool

4. If symptoms do not improve after taking the product for 5 or 6 days, discontinue use and bring this instruction to a consultation with a doctor, pharmacist, or registered seller. (There may be other potential diseases)

Dosage and Administration

Peel off the surface liner (film) and apply to the affected area once a day.

<Pre><Pre>cautions related to dosage and administration>

(1) Strictly follow the dosage and directions. (2) This product is intended to relieve symptoms such as pain or swelling. It is not intended to cure the causes of pain or swelling. Therefore, apply this product only when symptoms appear. (3) Do not use more than 4 sheets per day. (4) If you are sweating or the affected area gets wet, dry it off thoroughly before applying. (5) Do not warm the applied area with a heating device or electric blanket, etc. (6) For persons with sensitive skin, cut the sheet into a 1-2 cm square and apply it to a delicate area such as the inner arm for at least half a day. Confirm that no symptoms such as rashes, redness, itching, or irritations appear before use.

Ingredients and Quantities (per 100g plaster)

	ingredicites and edulitities (per 100g plaster)			
	Ingredient	Quantity	Effect	
	Loxoprofen sodium hydrate	5.67g (5 g as anhydride)	Suppresses prostaglandin, the source of inflammation and pain.	
	Nonanoic acid vanillylamide	0.015g	The warming comfort effect promotes blood circulation in the affected area and relieves pain.	

Additives: Styrene-isoprene-styrene-block copolymer (SIS), polyisobutylene, hydrogenated rosin glycerol ester, BHT, I-menthol, liquid paraffin, and 2 other ingredients.

[1g plaster per sheet (7cm×10cm)]

Indications

Lower-back pain, shoulder pain associated with stiff shoulders, joint pain, muscle pain, tendonitis (hand/wrist pains), elbow pain (tennis elbow, etc.), bruises, sprains

Storage and Handling Precautions

(1) Store in a cool place with little humidity and out of direct sunlight. (2) Keep out of reach of children. (3) Don't put it in a different container. (It can lead to misuse or a lowering of quality) (4) To maintain product quality, store unused sheets in the bag and seal it properly. (5) Please do not take medicine after the expiration date. It is recommended to use the product shortly after opening an aluminum bag.

If you have any questions about this product, please contact the store where you purchased it or at the following address:

Daiichi Sankyo Healthcare Co., Ltd. Customer Relations Office 3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan zip/postal code 103-8234 TEL: 0120-337-336

Reception Hours: 9:00 am -5:00 pm (closed Saturday, Sunday, and holidays)

https://www.daiichisankyo-hc.co.jp/

01@registered trademark Made in Japan

Relief System for Sufferers from Adverse Drug Reactions TEL0120 - 149 - 931

Daiichi Sankyo Healthcare Co., Ltd.

3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan

Manufacturer and seller

Lead Chemical Co., Ltd.

77-3, Himata, Toyama-City, Toyama, Japan