

For severe pain in low back and shoulder Cool type

MADE IN JAPAN

# ロキソニン®Sパップ

## Loxonin S Poultice

Topical anti-inflammatory medicine



Pharmacist's intervention required medicine

### Indications

Relief of low back pain, shoulder pain associated with stiff shoulder, joint pain, muscle pain, peritendinitis (pain in hands/wrists), elbow pain (e.g. tennis elbow), bruise, and sprain.

### Dosage and Administration

Once daily, peel off the liner (film) from the surface of poultice, and apply it to the affected area.

〈Precautions related to Dosage and Administration〉

- (1) Follow the Dosage and Administration strictly.
- (2) This product is not for curing the disease that is causing pain or swelling, but for merely treating symptoms such as pain and swelling. Use this product only when you have relevant symptoms.
- (3) Do not use more than two pieces of poultice per day.
- (4) When the affected area is sweaty or wet, wipe there off well before use.
- (5) For people having sensitive skin, verify before using this product that it does not cause any symptoms such as rash/redness, itchiness, and irritation by applying a small piece (approx. 1-2 cm square) of the poultice to the sensitive part on the inside of the arm for more than half a day.

### Ingredients and Quantities

In 100 g of paste

Ingredient	Quantity
Loxoprofen sodium hydrate	1.134g (1g as anhydrate)

Other ingredients : Peppermint oil, polysorbate 80, titanium oxide, tartaric acid, disodium edetate, glycerol, carmellose sodium, talc, dried aluminum hydroxide, crotamiton, partially neutralized polyacrylic acid, and other 2 ingredients

[1 piece of poultice (10cm × 14cm) contains 10g of paste]

## Precautions

### Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects may occur more readily.)

1. This product should not be used by the following persons:

- (1) Persons who have had allergic symptoms with this product or its ingredients.
- (2) Persons who have developed asthma after taking this product, other cold medicines, antipyretic analgesics or topical anti-inflammatory medicine.
- (3) Children under 15 years.

2. Do not apply to the following area.

- (1) Surrounding area of the eyes, and mucosal membrane or such.
- (2) Affected area with eczema, rash, and wound.
- (3) Affected area with athlete's foot and ringworm, or the area with suppuration.

3. Do not use other topical anti-inflammatory medicine while using this product.

4. Do not use this product continuously for two weeks or longer.

(This product should be used for temporary relief of pain. If the pain or other symptoms persist, stop using this product and consult a doctor .)

### Things to consult

1. The following persons should consult a doctor or pharmacist before use.

- (1) Persons who are treated by a doctor.
- (2) Persons who have had allergic symptoms with medicines or such.
- (3) Women who are pregnant or considered to be pregnant
- (4) Elderly people
- (5) Persons who have been diagnosed as the following conditions:  
Bronchial asthma