

For runny nose, stuffy nose, cough, sputum, sore throat and fever

MADE IN JAPAN

# 新ルル<sup>®</sup>AゴールドDX

## Shin Lulu A Gold DX

Common cold medicine

Designated Class 2 OTC medicine



### Indications

Relief of cold symptoms (runny nose, stuffy nose, cough, sputum, sore throat, fever, chills, headache, sneezing, joint pain, and muscle pain)

### Dosage and Administration

Take the following amount with cold or warm water.

Age	Amount per dose	Daily dose
Adults(15 years or over)	3 tablets	3 times. Take within 30 minutes after a meal if possible.
12 to 14 years	2 tablets	
Under 12 years	Do not use.	

<Precautions related to Dosage and Administration>

- (1) Follow the Dosage and Administration strictly.
- (2) Use for children of twelve years and over should be supervised by guardians.

### Ingredients and Quantities

This products is a white sugar-coated tablet. Nine tablets contain the following ingredients.

Ingredient	Quantity
Clemastine fumarate	1.34 mg (1 mg as clemastine)
Belladonna total alkaloids	0.3 mg
Bromhexine hydrochloride	12 mg
Tranexamic acid	420 mg
Acetaminophen	900 mg
<i>dl</i> -methylephedrine hydrochloride	60 mg
Dihydrocodeine phosphate	24 mg
Anhydrous caffeine	60 mg
Benfotiamine (vitamin B <sub>1</sub> derivative)	24 mg

Other ingredients: Cellulose, CMC-Ca, hydroxypropyl cellulose, magnesium stearate, hypromellose, macrogol, titanium oxide, refined sugar, polyvinyl alcohol (partially saponified), talc, calcium carbonate, carnauba wax

## Precautions

### Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects/accidents may occur more readily.)

1. This product should not be used by the following persons:
  - (1) Persons who have had allergic symptoms with this product or its ingredients.
  - (2) Persons who have developed asthma after taking this product, other cold medicines or antipyretic analgesics.
  - (3) Children under 12 years
2. Do not use any of the following medicines while taking this product:  
Other cold medicines, antipyretic analgesics, sedatives, antitussives, expectorants, oral medicines containing antihistamine (oral medicines for rhinitis, motion sickness medicines, anti-allergic medicines or such), gastrointestinal analgesic/ antispasmodic medicines, oral medicines containing tranexamic acid
3. Do not drive vehicles or operate machines after taking this product.  
(Symptoms such as sleepiness, blurred vision, or abnormal dazzling may occur.)
4. Women who are breastfeeding should not take this product or should avoid breastfeeding if take it.
5. Do not drink alcohol before or after taking this product.
6. Do not use this product continuously for a long term.

### Things to consult

1. The following persons should consult a doctor or pharmacist before use:
  - (1) Persons who are receiving medical treatment from a doctor or dentist.
  - (2) Women who are pregnant or suspect that they are pregnant.
  - (3) Elderly people.
  - (4) Persons who have had allergic symptoms with medicines or such.
  - (5) Persons who have the following symptoms:  
High fever, dysuria
  - (6) Persons who have been diagnosed as the following conditions:  
Thyroidal functional disorder, diabetes, heart disease, hypertension, liver disease, kidney disease, gastric/duodenal ulcer, glaucoma, persons who have a thrombosis (cerebral thrombosis, myocardial infarction, thrombophlebitis or such), persons who may develop a thrombosis, respiratory functional disorder, obstructive sleep apnea syndrome, obesity