

For sore throat, runny nose and fever

MADE IN JAPAN

新ルル-A錠s

Shin Lulu A Tablets s

Common cold medicine

Designated Class 2 OTC medicine



Indications

Relief of cold symptoms (runny nose, stuffy nose, sneezing, sore throat, cough, sputum, chills, fever, headache, joint pain, and muscle pain)

Dosage and Administration

Take the following amount with cold or warm water.

Age	Amount per dose	Daily dose
Adults (15 years or over)	3 tablets	3 times. Take within 30 minutes after a meal if possible.
<u>12 to 14 years</u>	2 tablets	
<u>Under 12 years</u>		<u>Do not use.</u>

<Precautions related to Dosage and Administration>

- (1) Follow the Dosage and Administration strictly.
- (2) Use for children of twelve years and over should be supervised by guardians.

Ingredients and Quantities

This product is a white sugar-coated tablet. Nine tablets contain the following ingredients.

Ingredient	Quantity
Acetaminophen	900 mg
Clemastine fumarate	1.34 mg
Dihydrocodeine phosphate	24 mg
Noscapine	48 mg
<i>dl</i> -methylephedrine hydrochloride	60 mg
Guaiacol sulfonic acid potassium	240 mg
Anhydrous caffeine	75 mg
Benfotiamine (vitamin B ₁ derivative)	24 mg

Other ingredients: Cellulose, magnesium aluminometasilicate, CMC-Ca, hydroxypropyl cellulose, magnesium stearate, gum arabic, titanium oxide, stearic acid, refined sugar, talc, hypromellose, polyvinyl acetal diethyl aminoacetate, polyvinyl alcohol (partially saponified), povidone, fumaric acid, carnauba wax

Precautions

Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects /accidents may occur more readily.)

1. This product should not be used by the following persons:
 - (1) Persons who have had allergic symptoms with this product or its ingredients.
 - (2) Persons who have developed asthma after taking this product, other cold medicines or antipyretic analgesics
2. Do not use any of the following medicines while taking this product.
Other cold medicines, antipyretic analgesics, sedatives, antitussives, expectorants, oral medicines containing antihistamine (oral medicines for rhinitis, motion sickness medicines, anti-allergic medicines or such)
3. Do not drive vehicles or operate machines after taking this product. (Sleepiness or such may occur.)
4. Women who are breastfeeding should not take this product or should avoid breastfeeding if take it.
5. Do not drink alcohol before and after taking this product.
6. Do not use this product continuously for a long term.

Things to consult

1. The following persons should consult a doctor or pharmacist before use.
 - (1) Persons who are receiving medical treatment from a doctor or dentist.
 - (2) Women who are pregnant or considered to be pregnant.
 - (3) Elderly people.
 - (4) Persons who have had allergic symptoms with medicines or such.
 - (5) Persons with the following symptoms:
High fever, dysuria
 - (6) Persons who have been diagnosed as the following conditions:
Thyroidal functional disorder, diabetes, heart disease, hypertension, liver disease, kidney disease, gastric/duodenal ulcer, glaucoma, respiratory functional disorder, obstructive sleep apnea syndrome, obesity