

Before using, be sure to read these written explanation.

Also please put it in a safe place, so that you can read them when necessary.

Works well on colds with severe coughing and phlegm

ルルアタック®CX

Lulu Attack®CX

Common cold medicine



Designated Class 2 OTC Medicine

Characteristics of Lulu Attack®CX

Common cold medicine that exhibits an excellent effect on colds with stubborn coughs and phlegm buildup through a combination of nine active ingredients including dihydrocodeine phosphate, noscapine, dl-methylephedrine hydrochloride, L-carbocisteine, and others.


1. Exhibits an excellent effect on stubborn coughs through two antitussive ingredients -- dihydrocodeine phosphate and noscapine – that work on the cough center.
2. Relieves stubborn coughs by expanding your bronchial tube, making it easier to breathe, through use of dl-methylephedrine hydrochloride.
3. Makes it easier to get rid of phlegm caught in your throat, one of the causes of coughs, by normalizing the mucous membrane through the expectorant ingredients in L-carbocisteine.
4. Exhibits an excellent effect on allergic symptoms of a cold (runny nose, stuffy nose) with d-chlorpheniramine maleate and by easing sore throat and fever through the anti-inflammatory ingredients in ibuprofen
5. Easy-to-carry with PTP packaging.

Indications

Relief of various symptoms of a common cold (cough, phlegm, sore throat, runny nose, stuffy nose, fever, chills, sneezing, headache, joint pain, muscle pain)

Administration and Dosage

Take the following amount with cold or warm water

Age	Amount per dose	Daily dose
Adult (15 years or over)	2 tablets 	3 times Take within 30 minutes after a meal if possible.
Under 15 years		Do not use

<Precautions related to Dosage and Administration>

(1) Follow the Dosage and Administration strictly.

(2) How to take out the tablets:

As shown in the diagram, use your finger to push the PTP sheet that contains the tablet. Take off the aluminum foil on the back and ingest the tablet. (If you mistakenly take the tablet with the wrapping on it, it could puncture the mucous membrane of the esophagus and lead to other unknown consequences.)



Ingredients and Quantities

This product is white film-coated tablet. Six tablets contain the following ingredients.

Ingredient	Quantity	Function
Ibuprofen	450mg	Reduces fever, relieves sore throat by working on 【prostaglandins】 that cause inflammation and pain.
Dihydrocodeine phosphate	24mg	Works on the cough center to relieve coughs.
Noscapine	48mg	
dl-methylephedrine hydrochloride	60mg	Expands the bronchial tubes, relieves coughs.
L-carbocisteine	750mg	Makes it easier to get rid of phlegm, one of the causes of coughs, by clearing the airways in the mucous membrane.
Glycyrrhizic acid	39mg	Eases inflammation of the mucous membranes in the nose and throat.
d-chlorpheniramine maleate	3.5mg	Suppresses allergic symptoms of a cold (runny nose, nasal congestion)
Anhydrous caffeine	75mg	Eases headaches
Benfotiamine (Vitamin B1 derivative)	25mg	Promotes recovery of physical strength depleted by a cold.

Additives: D-mannitol, cellulose, polyoxyethylene hydrogenated castor oil, croscarmellose sodium, hypromellose, hydroxypropyl cellulose, talc, silicic anhydride, polyvinyl alcohol (partial Saponified product), triethyl citrate, titanium oxide, carnauba wax

Precautions

Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects/accidents may occur more readily.)

1. This product should not be used by the following persons:
 - (1) Persons who have had allergic symptoms with this product or its ingredients.
 - (2) Persons who have developed asthma after taking this product, other cold medicines or antipyretic analgesics.
 - (3) Children younger than 15.
 - (4) Expectant women within 12 weeks of delivery date.
2. Do not use any of the following medicines while taking this product:
Other cold medicines, antipyretics, analgesics, sedatives, antitussives, expectorants, oral medicines containing antihistamine (oral medicines for rhinitis, motion sickness medicines, anti-allergic medicines or such)
3. Do not drive vehicles or operate machines after taking this product (Sleepiness or such may occur).
4. Women who are breastfeeding should not take this product or should avoid breastfeeding if take it.
5. Do not drink alcohol before or after taking this product.
6. Do not take this product for more than 5 days .

Things to consult

The following persons should consult a doctor or pharmacist before use.

- (1) Persons who are treated by a doctor or dentist.
- (2) Women who are pregnant or considered to be pregnant.
- (3) Elderly people.
- (4) Persons who have had allergic symptoms with medicines or such.
- (5) Persons who have the following symptoms: High fever, dysuria
- (6) Persons who have been diagnosed as the following conditions:
Thyroid functional disorder, diabetes, heart disease, hypertension, liver disease, kidney disease, glaucoma, systemic lupus erythematosus, mixed connective tissue disease, respiratory functional disorder, obstructive sleep apnea syndrome, obesity
- (7) Persons who experienced any of the following diseases: gastric/duodenal ulcer, ulcerative colitis, Crohn's disease

Precautions for storage and handling

- (1) Please store in a cool and dry place away from sunlight.
- (2) Please keep out of the reach of children.
- (3) Please do not transfer this product to a different container. (It may cause misuse and affect the quality.)
- (4) Please do not use after the expiration date printed on the label.

Please direct questions about this product to the store where you purchased it or to the below address:

Customer Relations Center
Dai-Ichi Sankyo Healthcare Co., Ltd.
3-14-10, Nihonbashi, Chuo-ku, Tokyo, 103-8234, Japan
TEL: 0120-337-336
Operation Hours: 9am-5pm (not including Sat., Sun. and holidays)

Please check at www.lulu.jp for more information about Lulu.

01® Registered trademark

Manufactured and Sold by
Dai-Ichi Sankyo Healthcare Co., Ltd.
3-14-10, Nihonbashi, Chuo-ku, Tokyo

Website <https://www.daiichisankyo-hc.co.jp/>

英語 (English) 2020.7 作成