

Works well on cold with sore throat, severe cold with a fever

MADE IN JAPAN

ルルアタック EX

Lulu Attack EX

Common cold medicine



Designated Class 2 OTC medicine

Indications

Relief of cold symptoms (sore throat, fever, chills, headache, runny nose, stuffy nose, sneezing, cough, sputum, joint pain, and muscle pain)

<<Use Lulu Attack EX for these>>



Dosage and Administration

Take the following amount with cold or warm water.

Age	Amount per dose	Daily dose
Adults (15 years or over)	2 tablets	3 times. Take within 30 minutes after a meal if possible.
Under 15 years		Do not use.

<Precautions related to Dosage and Administration>

(1) Follow the Dosage and Administration strictly.

Ingredients and Quantities

Six tablets of this product contain the following ingredients.

Ingredient	Quantity
Tranexamic acid	750 mg
Ibuprofen	450 mg
Clemastine fumarate	1.34 mg (1 mg as clemastine)
Bromhexine hydrochloride	12 mg
<i>dl</i> -methylephedrine hydrochloride	60 mg
Dihydrocodeine phosphate	24 mg
Thiamine nitrate (Vitamin B1 nitrate)	25 mg
Riboflavin (Vitamin B ₂)	12 mg

Other ingredients: Trehalose, cellulose, macrogol, CMC-Na, polyvinyl alcohol (partially saponified), anhydrous silica, hydrogenated oil, magnesium stearate, hypromellose, titanium oxide, carnauba wax

<Precautions related to Ingredients and Quantities>

The riboflavin (vitamin B₂) contained in this product may turn the urine yellow.

Precautions

Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects/accidents may occur more readily.)

1. This product should not be used by the following persons:
 - (1) Persons who have had allergic symptoms with this product or its ingredients.
 - (2) Persons who have developed asthma after taking this product, other cold medicines or antipyretic analgesics.
 - (3) Children under 15 years.
 - (4) Pregnant women who are scheduled to give birth within 12 weeks.
2. Do not use any of the following medicines while taking this product:
Other cold medicines, antipyretics, analgesics, sedatives, antitussives, expectorants, oral medicines containing antihistamine (oral medicines for rhinitis, motion sickness medicines, anti-allergic medicines or such), oral medicines containing tranexamic acid
3. Do not drive vehicles or operate machines after taking this product.(Sleepiness or such may occur.)
4. Women who are breastfeeding should not take this product or should avoid breastfeeding if take it.
5. Do not drink alcohol before or after taking this product.
6. Do not use for more than 5 days.

Things to consult

1. The following persons should consult a doctor or pharmacist before use:
 - (1) Persons who are receiving medical treatment from a doctor or dentist.
 - (2) Women who are pregnant or considered to be pregnant.
 - (3) Elderly people.
 - (4) Persons who have had allergic symptoms with medicines or such.
 - (5) Persons who have the following symptoms: High fever, dysuria
 - (6) Persons who have been diagnosed as the following conditions:
Thyroidal functional disorder, diabetes, heart disease, hypertension, liver disease, kidney disease, glaucoma, systemic lupus erythematosus, mixed connective tissue disease, persons who have a thrombosis (cerebral thrombosis, myocardial infarction, thrombophlebitis or such), persons who may develop a thrombosis, respiratory functional disorder, obstructive sleep apnea syndrome, obesity
 - (7) Persons who have had the following diseases:
Gastric/duodenal ulcer, ulcerative colitis, Crohn's disease