

Works well on severe cold with runny nose and stuffy nose

MADE IN JAPAN

ルルアタック[®] NX

Lulu Attack NX

Common cold medicine



Designated Class 2 OTC medicine

Indications

Relief of cold symptoms (runny nose, stuffy nose, sore throat, cough, sputum, fever, chills, sneezing, headache, joint pain, and muscle pain)

Dosage and Administration

Take the following amount with cold or warm water.

Age	Amount per dose	Daily dose
Adults (15 years or over)	2 tablets	3 times. Take within 30 minutes after a meal if possible.
Under 15 years		Do not use.

<Precautions related to Dosage and Administration>

(1) Follow the Dosage and Administration strictly.

Ingredients and Quantities

Six tablets contain the following ingredients.

Ingredient	Quantity
Ibuprofen	450 mg
Belladonna total alkaloids	0.3 mg
Clemastine fumarate	1.34 mg
Bromhexine hydrochloride	12 mg
Dihydrocodeine phosphate	24 mg
dl-methylephedrine hydrochloride	60 mg
Anhydrous caffeine	75 mg
Benfotiamine (vitamin B ₁ derivative)	25 mg
Riboflavin (vitamin B ₂)	12 mg

Other ingredients: Corn starch, lactose, cellulose, CMC-Na, hydroxypropyl cellulose, magnesium stearate, polyvinyl alcohol-acrylic acid-methyl methacrylate copolymer, talc, titanium oxide, hypromellose, carnauba wax

<Precautions related to Ingredients and Quantities>

The riboflavin (vitamin B₂) contained in this product may turn the urine yellow.

Precautions

Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects/accidents may occur more readily.)

1. This product should not be used by the following persons:
 - (1) Persons who have had allergic symptoms with this product or its ingredients.
 - (2) Persons who have developed asthma after taking this product, other cold medicines or antipyretic analgesics.
 - (3) Children under 15 years
 - (4) Pregnant women who are scheduled to give birth within 12 weeks.
2. Do not use any of the following medicines while taking this product.
Other cold medicines, antipyretics, analgesics, sedatives, antitussives, expectorants, oral medicines containing antihistamine (oral medicines for rhinitis, motion sickness medicines, anti-allergic medicines or such), gastrointestinal analgesic/antispasmodic medicines
3. Do not drive vehicles or operate machines after taking this product.
(Symptoms such as sleepiness, blurred vision or abnormal dazzling may occur.)
4. Women who are breastfeeding should not take this product or should avoid breastfeeding if take it.
5. Do not drink alcohol before and after taking this product.
6. Do not use for more than 5 days.

Things to consult

1. The following persons should consult a doctor or pharmacist before use.
 - (1) Persons who are receiving medical treatment from a doctor or dentist.
 - (2) Women who are pregnant or considered to be pregnant.
 - (3) Elderly people.
 - (4) Persons who have had allergic symptoms with medicines or such.
 - (5) Persons with the following symptoms:
High fever, dysuria
 - (6) Persons who have been diagnosed as the following conditions:
Thyroidal functional disorder, diabetes, heart disease, hypertension, liver disease, kidney disease, glaucoma, systemic lupus erythematosus, mixed connective tissue disease, respiratory functional disorder, obstructive sleep apnea syndrome, obesity
 - (7) Persons who have had the following diseases:
Gastric/duodenal ulcer, ulcerative colitis, Crohn's disease