

Inconspicuous transparent patch which avoid scratching

MADE IN JAPAN

マキロン®パッチエース

Makiron Patches Ace

Antipruritic and antiphlogistic patch

Designated class 2 OTC medicine



Indications

Itching due to insect bites

Dosage and Administration

Apply to the affected area 1 to 3 times a day.

<Precautions related to Dosage and Administration>

- (1) Follow the Dosage and Administration strictly.
- (2) Use for children should be supervised by guardians.
- (3) This product is for external use only.
- (4) When the affected area is sweating or wet, wipe away sufficiently before use.
- (5) If itching resolves after using this product for a few hours, please remove it.
- (6) Do not use at the same site for a long time.

Ingredients and Quantities

In 100 g (1.25 m²) of plaster of this product:

Ingredients	Quantity
Dexamethasone acetate	0.025 g
Diphenhydramine	1.00 g
Isopropyl methylphenol	1.00 g
Tocopherol acetate (acetate ester of vitamin E)	0.50 g

Additives: Diisopropyl adipate, methacrylate/ butylacrylate copolymer, BHT, *l*- menthol

This product contains dexamethasone acetate, an adrenocortical hormone (steroid).

Precautions

Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects may occur more readily.)

1. Do not use at the following sites:

- (1) Varicella (chickenpox), athlete's foot, ringworm or suppurative area
- (2) Area around the eyes, mucous membrane of the eyes and such
- (3) Wound, moist skin (scratch), erosion, eczema, eruption

2. This product should not be used on large areas of the face.

3. Do not use this product continuously for a long term.

Things to consult

1. The following persons should consult a doctor or pharmacist before use.

- (1) Persons who are receiving medical treatment from a doctor.
- (2) Women who are pregnant or considered to be pregnant.
- (3) Persons who have had allergic symptoms with medicines or such.
- (4) Persons whose affected areas are large.