

For swollen throat, sore throat, and stomatitis

MADE IN JAPAN

ペラック[®]T錠

Pelack T Tablets

Oropharyngeal medicine

Class 3 OTC medicine



Indications

Pharyngitis/tonsillitis (throat swelling and throat pain), stomatitis

Dosage and Administration

Take the following amount with cold or warm water.

Age	Amount per dose	Daily dose
Adults(15 years or over)	2 tablets	3 times. Take the tablets in the morning, at noon, and in the evening.
7 to 14 years	1 tablet	
Under 7 years	Do not use.	

<Precautions related to Dosage and Administration>

- (1) Follow the Dosage and Administration strictly.
- (2) Available to take either before or after meals.
- (3) Use for children aged 7 or over should be supervised by guardians.

Ingredients and Quantities

This product is a pale yellow film-coated tablet. Six tablets contain the following ingredients.

Ingredient	Quantity
Tranexamic acid	750 mg
Powdered extract of glycyrrhiza	198 mg (990 mg as crude drug)
Pyridoxine hydrochloride (Vitamin B ₆)	50 mg
Riboflavin (Vitamin B ₂)	12 mg
Sodium L-ascorbate (Vitamin C sodium)	500 mg

Other ingredients: Cellulose, silicic anhydride, polyvinyl alcohol (partially saponified), corn starch, magnesium stearate, hypromellose, titanium oxide, talc, propylene glycol, dimethylpolysiloxane, silicon dioxide, iron sesquioxide, carnauba wax

<Precautions related to Ingredients and Quantities>

- (1)The riboflavin (Vitamin B₂) contained in this product may turn the urine yellow.
 - (2)Taking this product may affect the results of laboratory tests performed on the urine or feces.
- If you are examined by doctors, tell them that you are taking a medicine containing Vitamin C.

Precautions

Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects may occur more readily.)

1. Do not use any of the following medicines while taking this product.
 - (1) Oral medicines which contain licorice or its main component, glycyrrhizinic acid.
(May cause edema, increase in blood pressure, or muscular disorders [myopathy] or such)
 - (2) Oral medicines containing tranexamic acid
(Oral medicines for rhinitis, cold medicines, antipyretic analgesics, antitussive and expectorant medicines or such)
2. Do not use this product continuously for a long term.

Things to consult

1. The following persons should consult a doctor or pharmacist before use.
 - (1) Persons who are receiving medical treatment from a doctor or dentist.
 - (2) Women who are pregnant or considered to be pregnant.
 - (3) Elderly people.
 - (4) Persons who have had allergic symptoms with medicines or such.
 - (5) Persons with the following symptom:
Edema
 - (6) Persons who have been diagnosed as the following conditions:
Hypertension, heart disease, kidney disease, persons who have a thrombosis (cerebral thrombosis, myocardial infarction, thrombophlebitis or such), persons who may develop a thrombosis