

<Pyrine medicine> Contains isopropylantipyrine

MADE IN JAPAN

プレコール®持続性カプセル

Precol Long-acting Capsule

Common cold medicine

Designated Class 2 OTC medicine



Indications

Relief of cold symptoms (sore throat, fever, runny nose, stuffy nose, cough, sputum, chills, headache, joint pain, muscle pain, and sneezing)

Dosage and Administration

Take the following amount with cold or warm water.

Age	Amount per dose	Daily dose
Adults (15 years or over)	2 capsules	twice (Take within 30 minutes after breakfast and dinner if possible.)
Under 15 years	Do not use.	

<Precautions related to Dosage and Administration>

(1) Follow the Dosage and Administration strictly.

Ingredients and Quantities

This product is a yellow and white capsule. Four capsules contain the following ingredients.

Ingredient	Quantity
Isopropylantipyrine (pyrine medicine)	300 mg
Acetaminophen	450 mg
Chlorpheniramine maleate	7.5 mg
Dihydrocodeine phosphate	12 mg
dl-methylephedrine hydrochloride	60 mg
Powdered extract of glycyrrhiza	118 mg (983 mg as crude drug)
Anhydrous caffeine	75 mg

Other ingredients: lactose, hydroxypropyl cellulose, corn starch, silicon dioxide, methacrylic acid copolymer L, stearic acid, talc, cellulose, CMC-Ca, gelatin, sodium lauryl sulfate

Precautions

Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects/accidents may occur more readily.)

1. This product should not be used by the following persons:
 - (1) Persons who have had allergic symptoms with this product or its ingredients.
 - (2) Persons who have developed asthma after taking this product, other cold medicines or antipyretic analgesics.
2. Do not use any of the following medicines while taking this product:
Other cold medicines, antipyretics, analgesics, sedatives, antitussives, expectorants, oral medicines containing antihistamine (oral medicines for rhinitis, motion sickness medicines, anti-allergic medicines or such)
3. Do not drive vehicles or operate machines after taking this product (Sleepiness or such may occur).
4. Women who are breastfeeding should not take this product or should avoid breastfeeding if take it.
5. Do not drink alcohol before or after taking this product.
6. Do not use this product continuously for a long term.

Things to consult

1. The following persons should consult a doctor or pharmacist before use.
 - (1) Persons who are treated by a doctor or dentist
 - (2) Women who are pregnant or considered to be pregnant
 - (3) Elderly people
 - (4) Persons who have had allergic symptoms with medicines or such.
 - (5) Persons who have the following symptoms: High fever, dysuria
 - (6) Persons who have been diagnosed as the following conditions:
Thyroidal functional disorder, diabetes, heart disease, hypertension, liver disease, kidney disease, gastric/duodenal ulcer, glaucoma, respiratory functional disorder, obstructive sleep apnea syndrome, obesity