

Apply directly to cure stomatitis and glossitis

MADE IN JAPAN

トラフル®軟膏

Traful Ointment

Ointment for stomatitis

Class 3 OTC medicine



Indications

Stomatitis, glossitis

Dosage and Administration

Apply a suitable amount to the affected area 2 to 4 times a day. Clean up the affected area before applying this product.

<Precautions related to Dosage and Administration>


- (1) Follow the Dosage and Administration strictly.
- (2) Use for children should be supervised by guardians.
- (3) Use only in the oral cavity.

How to use Traful Ointment

1. Rinse your mouth before using this product.
2. Apply the ointment so that it covers the affected area.
3. After applying the ointment, manage to avoid touching to the affected area.

< How to open the tube >

Invert the cap and press the inside protrusion firmly against the tip of the tube to open.



Ingredients and Quantities

This product contains the following ingredients per 100 g.

Ingredient	Quantity
Azulene sodium sulfonate hydrate	0.02 g
Glycyrrhetic acid	0.3 g
Allantoin	0.3 g
Cetylpyridinium chloride hydrate	0.1 g

Other ingredients: Glycerin, gelatinized hydrocarbon, sodium polyacrylate, magnesium aluminometasilicate, hypromellose, *l*-menthol, sodium saccharate

Precautions

Things to consult

1. The following persons should consult a doctor, dentist or pharmacist before use.
 - (1) Persons who are receiving medical treatment from a doctor or dentist.
 - (2) Persons who have had allergic symptoms with medicines or such.
 - (3) Persons who have wide range of inflammation inside the mouth.