For stomatitis and sore throat

トラフルウィックショット

Traful Quick Shot

Anti-inflammatory spray for oral cavity

Class 3 OTC medicine



Indications

Stomatitis,

Throat pain, sore throat, throat swelling, throat discomfort and hoarseness caused by inflammation of the throat.

Dosage and Administration

Several times a day, spray proper amount to the affected area.

<Precautions related to Dosage and Administration>

- (1) As the use of this product while breathing in may allow medicinal liquid to enter the bronchial tubes or lungs, utter a sound while spraying if the back of throat is targeted.
- (2) Keep the nozzle away from oral cavity while use.
- (3) Follow the Dosage and Administration strictly.
- (4) Use for children should be supervised by guardians.
- (5) Prevent the mist of spray from entering into the eyes. If it happens, rinse the eyes immediately with water or lukewarm water. If symptoms are severe, see an ophthalmologist.

[How to use the container]

After removing the cap, point the nozzle toward the affected area and pump to spray. When using for the first time, pump 4 or 5 times until the medicinal liquid comes out.

000

Ingredients and Quantities

Ingredients and quantities (per 100 g)

Ingredient	Quantity
Azulene sodium sulfonate hydrate (water-soluble azulene)	20 mg

Other ingredients: *I*-menthol, benzethonium chloride, glycerin, propylene glycol, polyoxyethylene hardened castor oil, carboxy vinyl polymers, sodium edetate, stevia extract, sodium chloride, pH modifier

Precautions

Things to consult

- 1. The following persons should consult a doctor or pharmacist before use.
 - (1) Persons who have had allergic symptoms with medicines or such.
 - (2) Persons with the following symptoms:
 - Severe sores or inflammation in the mouth