

Please read this instruction before use. Please keep this in a safe place so that you can read it when you need to.

Gastrointestinal Medicine

Class 2 OTC medicine

Fast effective relief of stomach heaviness, retching, and stomach discomfort.

第一三共胃腸薬錠剤s

Daiichi Sankyo Ichoyaku (Tablets) s

Product Details

This medicine product works to promote the relief of stomach heaviness, retching, and stomach discomfort as per the following:

- Aids digestion through the utilization of the fat digestion enzyme lipase AP12 and the digestive enzyme taka-diestase N1.
- Promotes the functioning of weakened stomachs and relieves stomach discomfort and loss of appetite with 6 types of stomachic ingredients medicinal herbs.
- Suppresses the cause of stomach pain with mallotus japonicus and Powdered glycyrrhiza for soothing irritation or inflammation of the gastric mucosa, and 3 types of antacids for restoring the balance of gastric acids.
- Does not contain sodium. Can be taken by persons concerned about sodium intake levels.



Precautions



Things to avoid

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

1. This product should not be taken by the following persons.

Persons receiving dialysis therapy.

2. Do not use continuously over extended periods.



Things to consult

1. Consult your doctor, pharmacist, or registered salesperson before taking if you meet any of the following conditions:

- (1) Persons who are receiving medical treatment from a doctor.
- (2) Persons who have had allergic symptoms with medicines or such.
- (3) Persons who have been diagnosed as the following condition:
Kidney disease

2. If the following symptoms occurs after taking this medicine, you could be experiencing side effects. In such case, stop taking it immediately and bring these instructions to a consultation with your doctor, pharmacist, or registered salesperson.

Related Part	Symptoms
Skin	Rash, redness, itching

3. The following symptoms may appear, but if any of those symptoms persist or increase, discontinue taking the product and bring these instructions to a doctor, pharmacist, or registered salesperson.

Constipation, diarrhea

4. If symptoms do not improve after 2 weeks, stop taking the product and bring these instructions to a doctor, pharmacist, or registered salesperson.

Indications

- Stomach heaviness, excessive eating or drinking, epigastric pressure, or loss of appetite.
- Heartburn, stomach pain, gastric hyperacidity, stomach heaviness, discomfort, or burping
- Poor digestion, promotes digestion, weak stomach, abdominal or stomach bloating
- Nausea (stomach retching, upset stomach caused by hangover/sickness from drinking), or vomiting



Stomach heaviness



Stomach retching





Stomach discomfort



Stomach pain

Dosage and Administration

Please take the following amount with water or lukewarm water.

Ages	Dosage	Doses/day
Age 15 and older	3 tablets 	3 times/day Take after meals
Age 11 to 14	2 tablets 	
Under 11 years	Do not use.	

<Precautions related to dosage and administration>

- (1) Strictly follow the dosage and directions.
- (2) For children aged 11 years and older, please follow the guidance of a guardian when taking this medicine.

Ingredients and Quantities

This medicine consists of light grey-brown to light brown uncoated tablets. 9 tablets contain the following ingredients.

Ingredient	Quantity	Effect
Taka-Diastase N1	150 mg	Digestive enzyme that can function across a wide pH range to aid in the digestion and absorption of nutrients.
Lipase AP12	60 mg	Fat digestive enzyme shown to be effective in aiding digestive function.
Mallotus bark extract	63 mg (504 mg as mallotus bark)	Suppresses excessive gastrointestinal tension, protects the gastric mucosa, and reduces stomach inflammation.
Powdered glycyrrhiza	150 mg	
Magnesium Aluminosilicate	720 mg	Long acting antacid effect (acid suppressant) works to protect the surface of the gastro mucosa. The synthetic hydrotalcite promotes a combined fast-acting effect.
Synthetic hydrotalcite	300 mg	
Magnesium hydroxide	600 mg	Fast-acting antacid effect (acid suppressant) helps to improve symptoms of hyperacidity including heartburn and burping.
Powdered phellodendron bark	105 mg	Aromas and flavors stimulate the taste buds and regulate the secretion of gastric and digestive juices and improve digestive function.
Powdered cinnamon bark	225 mg	
Powdered fennel	60 mg	
Powdered clove	30 mg	
Powdered ginger	75 mg	
I-Menthol	9 mg	

Additives: zanthoxylum fruit, potato starch, carmellose, magnesium stearate

Storage and Handling Precautions

- (1) Please tightly seal the bottle and store in a cool place with little humidity and out of direct sunlight.
- (2) Store out of reach of children.
- (3) Don't put in a different container. (It can lead to misuse or lowering of quality.)
- (4) Do not handle with wet hands. The surface may dissolve if wet and cause discoloration or unevenness of color. Do not return wet or moist tablets to the bottles as it may affect other tablets.
- (5) The filling in the bottle is to prevent damage to the product during transportation. Please discard after opening the bottle.
- (6) Do not use the product after the expiration date indicated on the outer packaging. Also, once the bottle is opened, use within 12 months.
- (7) Please write the date of opening the bottle on the "Opening date" section of the label.

If you have any questions about this product, please contact the store where you purchased it or at the following address:

Daiichi Sankyo Healthcare Co., Ltd. Customer Relations Office

3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan zip/postal code 103-8234

TEL: 0120-337-336

Reception Hours: 9:00 am -5:00 pm (closed Saturday, Sunday, and holidays)

<https://www.daiichisankyo-hc.co.jp/>



Manufacturer and seller

Daiichi Sankyo Healthcare Co., Ltd.

3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan