

Please read this instruction before use.

Please keep this in a safe place so that you can read it when you need to.

H2 blockers gastrointestinal drug

Heartburn due to gastric acid reflux

Fast-dissolving tablets that can be taken without water

ガスター-10 S錠 リバースコントロール

Gaster10 S Tablet Reverse Control

Class 1 OTC medicine

- If symptoms do not improve after 3 days of taking the product, stop taking it and consult your doctor, or pharmacist with this instruction.
- Do not take continuously for more than 2 weeks.
(Seek medical attention because serious gastrointestinal problems may be overlooked.)



Product Details

- This is a different type of gastrointestinal drug from acid-neutralizing gastrointestinal drugs. It controls excessive stomach acid and has an excellent effect on heartburn caused by stomach acid reflux.
- It can be taken before bedtime, on an empty stomach, or whenever symptoms such as heartburn, stomach pain, or upset stomach appear.
- Conveniently packaged in the PTP type for easy carrying.
- Fast-dissolving oral tablets allow for administration without water.

Precautions

Things to avoid

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

1. This product should not be taken by the following persons.
 - (1) Persons who have experienced allergic symptoms (e.g., rash/redness, itching, swelling of the throat, eyelids, lips, etc.) due to H2 blockers such as famotidine
 - (2) Persons who are receiving medical treatment for the following illnesses or receiving medicines at a medical institution
Blood diseases, kidney and liver diseases, heart diseases, stomach and duodenal diseases, immune system diseases such as asthma and rheumatism, steroids, antibiotics, anticancer drugs,azole antifungal drugs
(May cause leukopenia, thrombocytopenia, etc.)
(If you have kidney or liver disease, the excretion of the drug may be delayed, and its effects may be more pronounced.)
(If you have a heart disease such as myocardial infarction, valvular disease, or cardiomyopathy, pulse sagging with electrocardiographic abnormalities may occur.)
(Persons undergoing treatment for gastrointestinal or duodenal diseases are likely to be prescribed famotidine or similar drugs.
They should be careful not to duplicate dosages of the same medication.)
(Decreases absorption ofazole antifungal agents and reduces their effectiveness)
 - (3) Persons who have been diagnosed by a doctor with blood disorders such as low red blood cell count (anemia), low platelet count (difficulty in stopping bleeding or tendency to bleed easily), low white blood cell count.
(This product may trigger repeated blood disorders.)
 - (4) Phenylketonuria (This drug contains the L-phenylalanine compound)
 - (5) Children (under the age of 15) and the elderly people (the age of 80 and older)
 - (6) Women who are pregnant or considered to be pregnant
2. Do not take the following medications while taking this medicine.
Other gastrointestinal medicines
3. Women who are breastfeeding should not take this drug or should avoid breastfeeding when taking this medicine.

Things to Consult

1. The following persons should consult a doctor, or pharmacist before taking.
 - (1) Persons who are receiving medical treatment from a doctor or taking other medicines

(2) Persons who have had allergic symptoms with medicines or such.

(3) Elderly (65 and older)

(In general, the elderly may have reduced physiological functions)

(4) Persons with the following symptoms

Sore throat, cough, and high fever (Persons with these symptoms may be suspected of having a serious infection, and blood abnormalities such as decreased blood cell counts may be observed. If such symptoms are present before taking the drug, they may be aggravated by taking the drug and may cause delay in recognizing the side effects of the drug), unexplained weight loss, persistent abdominal pain (which may be caused by other diseases).

2. If any of the following symptoms appear after taking this medicine, you could be experiencing side effects. In such case, stop taking it immediately and consult a doctor, or pharmacist with this instruction.

Parts of the body affected	Symptoms
Skin	Rash/redness, itching, swelling
Circulatory system	Irregular pulse
Psycho-nervous system	Feeling of faintness, convulsions
Other	Abnormal physical symptoms such as feeling sick, lethargy, fever and sore throat

In rare cases, the following serious symptoms may occur. In such cases, seek medical attention immediately.

Name of symptoms	Symptoms
Shock (anaphylaxis)	Skin itching, hives, hoarse voice, sneezing, itchy throat, breathlessness, palpitations, and confusion of consciousness appear soon after taking this medicine.
Mucocutaneous-ocular Syndrome (Stevens-Johnson Syndrome) Toxic epidermal necrolysis	High fever, red eyes, eye discharge, sores on the lips, sore throat, and extensive skin rash/redness persist or rapidly worsen.
Rhabdomyolysis	Muscle pain in limbs, shoulders, and hips, numbness, weakness, stiffness, general lethargy, reddish brown urine are observed.
Liver dysfunction	Fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, general lethargy, and loss of appetite may occur.
Renal impairment	Fever, rash, decreased urine output, general swelling, general lethargy, joint pain (aching joints), and diarrhea may occur.
Interstitial pneumonia	Shortness of breath or difficulty breathing when climbing stairs or exerting oneself a little, dry cough, fever are observed, and these symptoms may appear suddenly or persist.
Blood disorders	Sore throat, fever, general lethargy, whiteness on the face or on the back of the eyelids, easy bleeding (bleeding gums, nosebleeds), blue bruises (color does not disappear when pressed), etc. are observed.

3. If you have accidentally taken more than the prescribed dosage, stop taking it immediately and consult your doctor, or pharmacist with this instruction.

4. If any of the following symptoms appear, and persist or increase after taking this medicine, discontinue taking the product immediately and consult a doctor, or pharmacist with this instruction.

Constipation, soft stool, diarrhea, dry mouth

Indications

Heartburn, stomachache, nausea, upset stomach (This medicine contains H2 blockers)

<Precautions related to indications>

Do not take this medicine for symptoms other than those listed in the Indications section.

Dosage and Administration

When symptoms of heartburn, stomachache, nausea, or retching appear, take the following dose dissolve in the mouth or take with water or lukewarm water.

Age	Dosage	Doses/day
Adult (Age 15 and older, under the age of 80)	1 tablet	Up to 2 times
Children (Under 15)		Do not take.
Elderly (80 and older)		Do not take.

- If symptoms persist for more than 8 hours after taking, take another 1 tablet.
- If symptoms subside, stop taking.
- If symptoms do not improve after 3 days, stop taking and consult a doctor, or pharmacist.
- Do not take continuously for more than 2 weeks.

<Precautions related to dosage and administration>

- (1) Strictly follow the dosage and administration guidelines.
- (2) Although the drug disintegrates easily in the mouth, it is not absorbed through the oral mucosa. After dissolving in the mouth, swallow the drug with saliva or with water or lukewarm water. As with ordinary tablets, taking the medicine directly with water or hot water does not change its effect.
- (3) Do not consume alcoholic beverages when taking this medicine. (Drugs are generally not to be taken with alcoholic beverages.)

[How to remove tablets]

As shown on the right, press firmly with your fingertip on the convex part of the PTP sheet containing the tablets to tear the aluminum foil on the back side, and take it out.
(If swallowed by mistake, it may cause an unexpected accident such as penetration of the esophageal mucosa.)



Ingredients and Quantities

This medicine is a white tablet, and each 1 tablet contains the following ingredients:

Ingredients	Quantity	Function
Famotidine	10 mg	Controls excessive stomach acid secretion.

Additives: Ethylcellulose, Cetanol, Sodium Lauryl Sulfate, Triacetin, Cyclodextrin, Flavor, l-Menthol, D-Mannitol, Aspartame(L-Phenylalanine Compound), Maltose Syrup Powder, Calcium Stearate

Storage and Handling Precautions

- (1) Store in a cool place with little humidity and out of direct sunlight.
- (2) Store out of reach of children.
- (3) Do not put in a different container. (It can lead to misuse or lowering of quality.)
- (4) Do not take medicine after the indicated expiration date.

This medication is not to be taken at set times, but rather when symptoms occur.

It is not affected by meals, so it can be taken before, after, or between meals.

One tablet at a time controls excessive stomach acid for about 8 hours, so if you take it twice a day, leave at least 8 hours between doses.

Maintaining gastrointestinal health

Binge eating and drinking, having an unbalanced diet, skipping meals are detrimental to gastrointestinal health. Avoid such dietary habits.

In addition, have regular health checkups.

If you have any questions about this product,
please contact the store where you purchased it or at the following address:

Daiichi Sankyo Healthcare Co., Ltd. Customer Relation Office

3-14-10, Nihonbashi, Chuo-ku, Tokyo, 103-8234, Japan

TEL: 0120-337-336

Reception Hours: 9:00 am - 5:00 pm (closed Saturdays, Sundays, and holidays)

<https://www.daiichisankyo-hc.co.jp/>



Manufacturer and seller

DAIICHI SANKYO HEALTHCARE CO., LTD.

3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan

English Ver. Created in Oct. 2024

This document is reference information about the products. Please confirm a document attached to a product definitely.