

Contains the same dosage of medical ingredients as the prescription drug

ロキソニン® S LOXONIN S Japanese Pharmacopoeia Loxoprofen Sodium Tablets		
Ingredients and Quantities (1 tablet)	Loxoprofen sodium hydrate	68.1 mg (60 mg as anhydrate)
Additives	Hydroxypropylcellulose, magnesium stearate, lactose monohydrate, red ferric oxide	

Some tablets may have red spots due to additives.

Added ingredients that are gentle on the stomach

ロキソニン® S プラス LOXONIN S PLUS		
Ingredients and Quantities (1 tablet)	Loxoprofen sodium hydrate	68.1 mg (60 mg as anhydrate)
	Magnesium oxide	33.3 mg
Additives	Lactose, cellulose, hydroxypropylcellulose, croscarmellose sodium, calcium stearate, hypromellose, titanium oxide, red ferric oxide, carnauba wax	

The tablet quickly dissolves after ingestion

ロキソニン® S クイック LOXONIN S QUICK		
Ingredients and Quantities (1 tablet)	Loxoprofen sodium hydrate	68.1 mg (60 mg as anhydrate)
	Magnesium aluminometasilicate	100 mg
Additives	Anhydrous dibasic calcium phosphate, lactose, croscarmellose sodium, hydroxypropylcellulose, magnesium stearate	

Formulated for headache relief

ロキソニン® S プレミアム LOXONIN S PREMIUM		
Ingredients and Quantities (2 tablets)	Loxoprofen sodium hydrate	68.1 mg (60 mg as anhydrate)
	Allylisopropylacetylurea	60 mg
	Anhydrous caffeine	50 mg
	Magnesium aluminometasilicate	100 mg
Additives	Lactose, cellulose, hydroxypropylcellulose, croscarmellose sodium, magnesium stearate, hypromellose, titanium oxide, talc, red ferric oxide, carnauba wax	

Focusing on the mechanism of menstrual pain

ロキソニン® S プレミアムファイン LOXONIN S PREMIUM fine		
Ingredients and Quantities (2 tablets)	Loxoprofen sodium hydrate	68.1 mg (60 mg as anhydrate)
	Paeonia dry extract	36 mg (252 mg as whole crude drug)
	Hesperidin	30 mg
	Magnesium aluminometasilicate	100 mg
Additives	D-Mannitol, cellulose, croscarmellose sodium, hydroxypropylcellulose, magnesium stearate, hypromellose, titanium oxide, macrogol, red ferric oxide, carnauba wax	

Quick relief from shoulder and lower back pain

ロキソニン® PROフィジカル LOXONIN PRO Physical		
Ingredients and Quantities (2 tablets)	Loxoprofen sodium hydrate	68.1 mg (60 mg as anhydrate)
	Paeonia dry extract	36 mg (252 mg as whole crude drug)
	Glycyrrhiza dry extract	50.4 mg (252 mg as whole crude drug)
	Magnesium oxide	33.3 mg
Additives	Lactose, partly pregelatinized starch, anhydrous silicic acid, croscarmellose sodium, carmellose calcium, calcium stearate, hypromellose, titanium oxide, macrogol, red ferric oxide, carnauba wax	

What effect does this medicine have?

- This medicine is an antipyretic analgesic that belongs to a group called non-steroidal anti-inflammatory drugs (NSAIDs).
- This medicine suppresses the production of "prostaglandins", the substances that cause pain, inflammation, and fever, thereby relieving pain and reducing fever.
- This medicine can be taken for the following purposes:
 - Pain relief: for headache, menstrual pain, toothache, pain after tooth extraction, sore throat, back pain, joint pain, neuralgia, muscle pain, stiff shoulder, earache, bruise, fracture, sprain, trauma pain
 - Fever reduction during chills and fever

What should be confirmed before purchasing this medicine?

Please check the following items to make sure that you can take this product.

It is known that side effects (adverse reactions) may include stomach/duodenal ulcers, induction of severe asthma attacks, rashes, and swelling.

For more information, please see [What caution should be taken when using this medicine?].

Do not take this medicine if:

- You have experienced allergic reactions to this medicine or its components in the past.
- You have developed asthma after taking this or other antipyretic analgesics, or cold medicines.
 - (If you have experienced asthma after taking antipyretic analgesics or cold medicines containing loxoprofen sodium hydrate, aspirin (acetylsalicylic acid), aspirin aluminum, ethenzamide, ibuprofen, acetaminophen, or isopropylantipyrine.)
- You are a child under the age of 15 years old.
- You are undergoing treatment for the following conditions at medical institutions:
 - Stomach/duodenal ulcers ●liver disease ●kidney disease ●heart disease
- Your doctor has diagnosed you with any of the following blood disorders:
 - Low red blood cell count (anemia)
 - Low platelet count (bleeds easily, difficulty stopping bleeding)
 - Low white blood cell count
- You are pregnant and due to give birth within 12 weeks

This medicine should be taken with caution in the following cases. Consult with a doctor, dentist or pharmacist before purchasing this medicine if:

- You are receiving medical or dental treatment.
- You are pregnant or might be pregnant (excluding pregnant women within 12 weeks of the expected delivery date).
- You are breastfeeding.
- You are elderly (65 years or older)
- You have experienced allergic reactions to medications before.
- You have been diagnosed with any of the following conditions:
 - Bronchial asthma ●ulcerative colitis ●Crohn's disease
 - systemic lupus erythematosus ●mixed connective tissue disease
- You have a history of any of the following diseases:
 - Stomach/duodenal ulcers ●liver disease ●kidney disease ●blood disease

What should be confirmed before using this medicine?

- This medicine is not intended to treat the underlying disease causing pain and fever; other, it is intended to suppress the symptoms that are occurring.
 - Therefore, **it should only be taken when symptoms are present.**

- If symptoms do not improve after 1 or 2 doses, stop taking the medicine, as this may indicate another condition and bring these instructions to a consultation with your doctor, dentist, or pharmacist.

- Do not use this medicine continuously for a long period of time. (If pain or other symptoms recur after 3 to 5 days of use, stop taking the medicine and seek medical attention.)**

- Do not take any of the following medicines while taking this medicine.
 - Other antipyretic analgesics, cold remedies containing aspirin (acetylsalicylic acid), aspirin aluminum, ethenzamide, ibuprofen, acetaminophen and isopropylantipyrine, and sedatives containing allylisopropylacetylurea and bromovalerylurea, **motion sickness medication [LOXONIN S PREMIUM only]**

- Do not drive a vehicle or operate machinery after taking this medicine. (Drowsiness and other symptoms may occur.) [LOXONIN S PREMIUM only]**

- Do not consume alcohol before or after taking this medicine.
 - [Do not take the medicine while consuming alcohol.]

How to use this medicine?

- Dosage and Administration**
 - For adults (15 years or older), should take the following dosage with water or warm water, preferably not on an empty stomach, when symptoms appear. The usual dosage **is up to 2 doses daily.**
 - If pain or other symptoms recur,** a third dose may be taken. The interval between doses should be at least 4 hours.

LOXONIN S	LOXONIN S PLUS	LOXONIN S QUICK
1 tablet per dose	1 tablet per dose	1 tablet per dose
LOXONIN S PREMIUM	LOXONIN S PREMIUM fine	LOXONIN S PRO Physical
2 tablets per dose	2 tablets per dose	2 tablets per dose

* **Be sure to read and strictly follow the enclosed instructions for dosage and administration when taking this product.**

- How should this medicine be taken?
 - Take it with a cup of water or warm water.
- What to do if you take too much of this medicine (overdose)?
 - In case of accidental overdose, consult a doctor or pharmacist.

What precautions should be taken when using this medicine?

- After taking antipyretic analgesics such as this medication, you may experience symptoms such as an excessive decrease in body temperature, weakness (lack of strength), and coldness of the limbs (cold hands and feet). If you experience any of these symptoms, stop taking the medicine immediately and bring these instructions to a consultation with your doctor or pharmacist.



- If you experience peptic ulcers (upset stomach, heartburn, back pain) or swelling after taking this medicine, stop taking it immediately and bring these instructions to a consultation with your doctor or pharmacist. In addition, severe symptoms such as gastrointestinal bleeding (vomiting blood, nausea and vomiting, abdominal pain, black tarry stools, bloody stools, etc.), gastrointestinal perforation (perforation of the gastrointestinal tract; nausea, vomiting, severe abdominal pain, etc.), and narrowing or obstruction of the small or large intestine (nausea, vomiting, abdominal pain, abdominal distention, etc.) may occur in rare cases. Seek medical attention immediately if you experience these symptoms.

What are the possible side effects?

- (1) If you experience any of the following symptoms after taking this medicine, they may indicate side effects. Discontinue use immediately and consult with a doctor, or pharmacist with the instruction leaflet.

Related Part	Symptoms
Skin	Rash, redness, itching
Digestive system	Abdominal pain, gastric discomfort, loss of appetite, nausea and vomiting, abdominal distention, heartburn, mouth ulcers, indigestion
Cardiovascular system	Increased blood pressure, palpitations
Neuropsychiatric system	Drowsiness, numbness, dizziness, headache
Other	Chest pain, fatigue, facial hot flashes, fever, anemia, hematuria

- (2) The following is a list of rare side effects that require special attention, along with their main subjective symptoms. If you experience a side effect, it is common for several of the main symptoms listed for that side effect to appear at the same time. In such cases, stop taking the medicine immediately and seek medical attention.

Symptom Names	Symptoms
Shock (anaphylaxis)	Symptoms such as itchy skin, urticaria, hoarseness, sneezing, itchy throat, difficulty breathing, palpitations, and clouding of consciousness may occur immediately after taking this medicine.
Blood disorders	Sore throat, fever, general lethargy, pallor face or eyelids, easy bleeding (bleeding gums, nosebleeds), blue bruises (bruises that do not fade when pressed) are observed.
Mucocutaneous ocular syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, erythema multiforme, acute generalized exanthematous pustulosis	High fever, eye redness, eye discharge, lip erosion, sore throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.
Kidney disorders	Symptoms such as fever, rash, reduced urinary output, full-body swelling, general malaise, joint pain and diarrhea, etc. may occur.
Congestive heart failure	General lethargy, palpitations, shortness of breath, chest discomfort, chest pain, dizziness, fainting, etc.
Myocardial infarction	Tight, constricting chest pain, difficulty breathing, and cold sweats may occur.
Cerebrovascular disorders	Symptoms such as reduced or loss of consciousness, difficulty moving the limbs on one side of the body, headache, vomiting, dizziness, slurred speech, and difficulty forming thoughts into words may occur suddenly.
Interstitial pneumonia	Shortness and/or difficulty breathing when climbing stairs or straining yourself slightly, sudden or persistent symptoms like dry cough, fever, etc.
Liver dysfunction	Symptoms such as fever, itching, rash, jaundice (skin and whites of the eyes turn yellow), brown urine, general malaise, loss of appetite, etc. may occur.
Rhabdomyolysis	Muscle aches in the limbs, shoulders, and hips; numbness, weakness, and stiffness in the limbs, general lethargy, and reddish-brown urine.
Aseptic meningitis	Symptoms such as severe headache accompanied by a tight sensation in the neck, fever, nausea/vomiting, etc. may occur. (These symptoms have frequently been reported particularly in individuals receiving treatment for systemic lupus

	erythematous or mixed connective tissue disease.)
Asthma	Wheezing or hissing when breathing, and difficulty breathing, etc. may occur.

The subjective symptoms listed in the [What precautions should be taken when using this medicine?] are sorted in the following list section above according to the body parts affected. If you notice any of these symptoms, please refer to the table in section above (2) "Side effects to especially watch out for" above.

Parts	Subjective Symptoms
Whole body	Lower than normal body temperature, lack of strength, general swelling, oedema, fever, general lethargy, high fever, drowsiness, muscle stiffness, cold sweats.
Head	Dizziness, severe headache with neck tightness, headache, fainting, decreased or total loss of consciousness.
Face	Face turns pale, nosebleeds, hot flashes
Eyes	Whitish underneath the eyelids, yellowing of the whites of the eyes, redness of the eyes, eye discharge
Ears	Tinnitus
Mouth and throat	Vomiting blood, sore throat, bleeding gums, nausea, vomiting, dry cough (cough without phlegm), mouth ulcers, sores on lips and in mouth, hoarse voice, sneezing, itchy throat, slurred speech, sudden difficulty speaking.
Chest	Heartburn, breathlessness, palpitations, shortness of breath, chest discomfort, clenching chest pain, chest pain, difficulty breathing, asthma (Symptoms, such as wheezing or hissing when breathing)
Abdomen	Upset stomach, abdominal pain, stomach discomfort, loss of appetite, abdominal distention, indigestion, severe abdominal pain
Back	Back pain, muscle pain in shoulders, lower back, etc.
Hands/Feet	Cold hands and feet, numbness, joint pain (pain in the joints), muscle pain, difficulty moving limbs on one side
Skin	Hives, blue bruises, rash, redness, widespread rash/redness, blisters on red areas of the skin, yellowing of the skin, itching
Feces	Black tarry stools, bloody stools, diarrhea
Urine	Decrease in urine volume, hematuria, brown urine, reddish brown urine
Other	Increased blood pressure, bleeding easily

- (3) The following symptoms may occur after taking this medicine. If these symptoms persist or worsen, stop taking the medicine, and bring the instructions to a consultation with your doctor, or pharmacist.

Dry mouth, constipation, diarrhea

Other

- How should this medicine be stored?
 - Store in a cool and dry place and out of direct sunlight.
 - Keep out of reach of children.
 - Do not transfer and keep in a different container. (This can lead to misuse or a decrease in quality.)

[LOXONIN S, LOXONIN S QUICK, LOXONIN PRO Physical]

- Please do not use the product after the indicated expiration date.

[LOXONIN S PLUS, LOXONIN S PREMIUM, LOXONIN S PREMIUM fine]

- Please do not use the product after the indicated expiration date. Also, once the aluminum bag is opened, use it within 12 months.
- Write the date you opened the aluminum bag in the "Date Opened" field on the box.

Who should be contacted about this medicine?

- For any general questions, please contact us at the address below.

Daiichi Sankyo Healthcare Co., Ltd.

(<https://www.daiichisankyo-hc.co.jp/>)

Customer Relations Office

3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan zip/postal code 103-8234

TEL: 0120-337-336

Reception Hours: 9:00 am -5:00 pm (Closed on Saturdays, Sundays, national holidays, and other days the company is closed)

- If you have more questions about symptoms, side effects, etc., please contact your doctor, dentist, or pharmacist. ©Registered Trademarks