

Analgesic and anti-inflammatory drug

Class 2 OTC Medicine

(2 patches x 10 sets)

LOXONIN S Tape mini 20 patches

(2 patches x 10 sets x 1 bag)

Before using this medicine, be sure to read the written instructions on this box. Also please keep it in a safe place so that you can read them when necessary.

Features

Formulated with loxoprofen sodium hydrate for effective analgesic and anti-inflammatory action.

Absorbed through the skin, converts to active form, and penetrates directly to the source of pain for effective relief.

Apply once daily. Comfortable to wear, with excellent elasticity, stays securely in place and is easy to reapply.

Scan for Product Info



Precautions

Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects may occur more readily.)

- Do not use this product if:
 - You have experienced allergic reactions to this product or its components in the past.
 - You have experienced asthma after using this product or other antipyretic analgesics, cold medicines, or topical analgesic and anti-inflammatory drugs.
 - You are a Child under 15 years of age.
- Do not use this product on:
 - The area around eyes or on mucous membranes.
 - Areas with eczema, rash, or wounds.
 - Areas affected by athlete's foot, ringworm, or infected wounds.
- Do not use other topical analgesic and anti-inflammatory products while using this product.
- Do not use this for more than 2 consecutive weeks. (This product temporarily relieves pain. If symptoms such as pain persist, discontinue use and consult a doctor.)

- If the following symptoms occur after use, they may indicate side effects. Stop using this product immediately and consult a doctor, pharmacist, or registered seller with this box.

Related Part	Symptoms
Skin	Rash/redness, itching, hives, stinging sensation, skin irritation, blisters, bruises, hyperpigmentation
Digestive system	Stomach discomfort, epigastric pain
Other	Swelling

In rare cases, the following severe symptoms may occur. In such cases, seek medical attention immediately.

Name of symptom	Symptoms
Shock (anaphylaxis)	Symptoms such as skin itching, urticaria, hoarseness, sneezing, throat itching, difficulty breathing, palpitations, or clouding of consciousness may occur immediately after use.

- The following symptoms may occur after use. If these symptoms persist or worsen, stop use and consult a doctor, pharmacist, or registered seller with this box. Diarrhea/loose stool
- If your symptoms do not improve after 5-6 days of use, discontinue use and consult a doctor, pharmacist, or registered seller with this box. (There may be other conditions.)



Things to consult

- Consult a doctor, pharmacist or registered seller before use this medicine if:
 - You are receiving medical treatment.
 - You have experienced allergic reactions to medications or other substances.
 - You are pregnant or might be pregnant.
 - You are elderly.
 - You have been diagnosed with: Bronchial asthma

Indications

Lower back pain, shoulder pain associated with stiff shoulders, joint pain, muscle pain, tendinitis (Hand/finger pain), elbow pain (e.g., tennis elbow), bruises, sprains

Dosage and Administration

Peel off the liner (film) from the surface and apply to the affected area once daily.

<Precautions related to Dosage and Administration>

- Strictly follow the dosage and administration instructions. (2) This product is intended solely for symptomatic relief of pain, swelling, and related symptoms, and does not address the underlying cause. It should only be used when symptoms are present. (3) Do not use more than 8 patches per day. (4) If the affected area is sweaty or wet, wipe it dry before use. (5) If you have sensitive skin, apply the product on a small area about 1-2 cm square on the inner arm where the skin is particularly sensitive, for at least half a day before full use, and ensure that no rash, redness, itching, or other irritation occurs.

Additives: Styrene-Isoprene-Styrene Block Copolymer, Polyisobutylene, Hydrogenated Rosin Glycerol Ester, BHT, I-Menthol, Liquid Paraffin, and 2 other ingredients **[Plaster mass per patch (5 cm x 7 cm): 0.5 g]**

Storage and Handling Precautions

- Store away from direct sunlight, in a cool, dry place.
- Keep out of reach of children.
- Do not transfer to a different container. (This may lead to misuse or quality deterioration.)
- To maintain quality, place any unused patches in the bag after opening and store with the interlocking zipper tightly closed.
- Do not use the product after the indicated expiration date. Use promptly after opening the aluminum bag.

If you have any questions about this product, please contact the store where you purchased it or at the following address:

Daiichi Sankyo Healthcare Co., Ltd. Customer Relations Office
3-14-10 Nihonbashi, Chuo-ku, Tokyo 103-8234
Phone: 0120-337-336 Reception Hours: 9:00 AM to 5:00 PM (excluding Saturdays, Sundays, and holidays)

<https://www.daiichisankyo-hc.co.jp/>
01@ Registered Trademarks

Sells Company
DAIICHI SANKYO HEALTHCARE CO., LTD.
3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan

Manufacturer and Marketing
Authorization Holder
Lead Chemical Co., Ltd.
77-3 Himata, Toyama City, Toyama

Relief System for Sufferers from
Adverse Drug Reactions
Phone: 0120-149-931

MADE IN JAPAN



4 987107 676955

Ingredients and Amount (per 100 g of ointment)

Ingredient	Amount	Function
Loxoprofen sodium hydrate	5.67 g (5 g as anhydrate)	Suppresses prostaglandins, which are the source of inflammation and pain.